



Directions for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no warranty, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from use in combination with other materials.

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
HARMFUL IF SWALLOWED

AVOID INHALATION OF VAPORS AND SPRAY MIST

NET CONTENTS 30 GALS.

# FLOW

## ACTIVE INGREDIENT:

By Weight

Petroleum Oil ..... 82.0%

INERT INGREDIENTS ..... 18.0%

Unsuifonated Residue, Petroleum Oil,  
Minimum ..... 92.0%

California Dept. of Agriculture Classification: Medium

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

**LEMONS:** Red Scale, Yellow Scale, Black Scale, Purple Scale, Soft Brown Scale, Citricola Scale, Citrus Red Mite, Citrus Bud Mite, Citrus Flat Mite, Two Spotted Mite. Use 1.7 to 2.0 gallons Flowable 75 per 100 gallons of water. Apply 400-3000 gallons of dilute spray per acre in thorough coverage. In Southern California, April and May and/or Sept. through Dec. In Central California, August and September.

## VALENCIA ORANGE OTHER CITRUS

Red Scale, Yellow Scale, Soft Brown Scale, Citrus Bud Mite. Use 1.7 to 2.0 gallons of water. Apply 400-3000 gallons of dilute spray per acre using thorough coverage. In areas, August and September. Do not use except on Navel.

For severe infestation, the addition of 1/2 cup of Rotolene before picking) is advisable. For one of the above spray solution ROTOLINE Aphidicide (at least 1 day before picking).



**LEFFINGWELL**  
**BREAST**

EPA REG. NO. 1148 30

GALS.

PRODUCT No. 3023

# LOWABLE 75

By Weight

82.0%

18.0%

um Oil,

92.0%

Classification: Medium

ON

OF CHILDREN

ale, Black Scale, Pur-  
Citricola Scale, Citrus  
s Flat Mite, Two Spot-  
s Flowable 75 per 100  
000 gallons of dilute  
rage. In Southern Cali-  
Sept. through Dec. In  
eptember.

## VALENCIA ORANGES, NAVEL ORANGES, GRAPEFRUIT, OTHER CITRUS VARIETIES:

Red Scale, Yellow Scale, Black Scale, Purple Scale, Soft Brown Scale, Citricola Scale, Citrus Red Mite, Citrus Bud Mite, Citrus Flat Mite, Two Spotted Mite — Use 1.7 to 2.0 gallons of Flowable 75 per 100 gallons of water. Apply 400-3000 gallons of finished spray per acre using thorough coverage. In Southern California areas, August only. In Central California, July and August except on Navel oranges July to August 15.

For severe infestations of Red, Yellow, or Purple Scale, the addition of 0.4 lb. Sevin 80 WP (at least 5 days before picking) or 1 lb. Malathion 25WP (at least 7 days before picking) per 100 gallons of spray solution is advisable. For severe Black Scale infestation, use one of the above or 1 1/4 lb. TOX-R™ per 100 gallons of spray solution (2-22 lbs TOX-R per acre) or 1 pint ROTOL™ Aphis Spray per 100 gallons of spray solution (at least 1 day before picking). Apply as soon as prac-

ticable following completion of major hatch. The rotenone treatment is suggested for emergency use when treatment has been delayed and larger stages are present.

## WARNING

Being an oil spray, application should NOT be made:

1. When the soil (especially the subsoil) is relatively dry.
2. Immediately prior to, during or following an unusually cold weather period.
3. When the temperature is above 95 degrees Fahrenheit.
4. Do not use combinations with Captan, Dinitro, Dichlone, Glyodin, Karathane, Sulfur, Lime Sulfur, Methoxychlor, or 60 days before or until 90 days after Sulfur, Captan, Phaltan products.
5. When used as a sticker-spreader, do not apply over fruit of apricots, cherries, apples or pears in excess of 0.2%.

Do not reuse empty drum. Return to manufacturer.

EFFINGWELL CHEMICAL COMPANY  
BREA, CALIFORNIA



EPA EST. NO. 1148-CA-1