

TOX-R™**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS****WARNING**

May be fatal if swallowed. May cause eye injury. Causes skin irritation. Avoid inhalation of dust. Do not get in eyes, on skin or on clothing. Wear protective goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

COMPOSITION

Active Ingredient: (% by weight)	
Rotenone	4.0%
Other Cube Resins	7.0%
Inert Ingredients:	89.0%
Total:	100.0%

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

KEEP OUT OF REACH OF CHILDREN**WARNING****STATEMENT OF PRACTICAL TREATMENT**

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites or gelatin solution. If not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.
IF IN EYES: Wash with plenty of clear water for at least 15 minutes. Call a physician.
IF ON SKIN: Wash with plenty of soap and water for at least 15 minutes. Call a physician if irritation persists. Remove and wash contaminated clothing before reuse.
IF INHALED: and abnormal symptoms develop, see a physician.

SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT.

EPA REG. NO. 1148-41
EPA EST. NO. 7874-CA-1

©Copyright 1987 Unifroyal Chemical Company, Inc.
NET CONTENTS:

001
12-15-89

ACCEPTED

MAY 24 1990

LEFFINGWELL

111 SOUTH BERRY STREET/P.O. BOX 1880, BREA, CA 92621

UNIFROYAL

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing unless sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"WARNING" Area to be treated with ROTENONE on (date of application). Do not enter without appropriate protective clothing until 24 hours after application.

DIRECTIONS FOR USE OF TOX-R

Utilizing the spray information provided, prepare the tank mix by adding the indicated amount of TOX-R to agitated water (unless otherwise indicated) containing a spreader at a rate of 1½ pints per 100 gals. Spray plants or trees lightly but thoroughly; cover all parts of plant or trees, particularly undersides of leaves. Do not apply if rain is expected shortly after application.

CROP	INSECTS	WHEN TO TREAT	AMOUNT
Citrus crops including Oranges, Lemons, Grapefruit, Tangerines, Kumquats (at least one day before harvest)	Aphids, leafroller, orange tortrix, red spider mite	When insects first appear. Spray as necessary.	With light medium oil emulsions (Fifinex only) Use 1/4-1/2 lb. TOX-R with 0.5-1 gal. of oil per 100 gals. water. Best results are obtained if the TOX-R is mixed with the oil and allowed to soak for a few minutes (between tanks) before adding to the spray tank. Apply 300-500 gals. of finished spray per acre (1 1/4-4 lbs. TOX-R per acre) as needed using outside coverage. With ROTO-FLO 50R (petroleum oil) (all varieties) Use 1/4-1/2 lb. TOX-R with 1/4-1 gal. ROTO-FLO 50R per 100 gals. water. results are improved if the TOX-R and oil are premixed and allowed to soak for a few minutes (between tanks) before adding to the spray tank. Apply 300-500 gals. of finished spray per acre (1/4-2 1/2 lbs. TOX-R per acre) as needed using outside coverage. With Men-oil spray Use 1 1/4-1 1/2 lbs. TOX-R with 1 1/2 pts. of a spreader-sticker per 100 gals. of water. Mix TOX-R in spreader and let soak between tanks. Apply 300-500 gals. finished spray per acre (3 1/4-7 1/2 lbs. TOX-R per acre) in outside coverage.
	Black, Brown, Citricole, and other soft scale insects	When insects first appear. Spray as necessary.	With light or medium grade or Narrow Range oils. (all varieties) Use 1/4-1/2 lb. TOX-R with 1.6-1.8 gals. emulsive (2 gals. Emulsion) Light Medium or Medium Oil or 1.2-1.4 gals. UNI-PAN™ per 100 gals. of water. Best results are obtained if TOX-R and oil are pre-mixed and allowed to soak for a few minutes before adding to spray tank. Apply 400-500 gals. of finished spray per acre (2-22 lbs. TOX-R per acre) using thorough coverage. This is suggested for emergency control of Black scale when treatment has been delayed and larger stages are present. Apply only during times when oil sprays are recommended.
Cranberry	cranberry spittle insect	When frothy spittle masses appear. Repeat as needed.	7 to 15 lbs. in 100 gals. per acre.
	fruit worms	When new growth is 2 to 4 inches long. Repeat at 10 day intervals.	
Apples, Pears	aphid, leafhopper, tent caterpillar, European apple sawfly, Japanese beetle, pear psylla, red spider mite, rapist bagworm	Start to spray when green leaf tips appear. Repeat at 10 day intervals until bloom, and at 14 day intervals after bloom.	10 to 20 lb. in 100 gal. per acre.
	codling moth, apple maggot	apply at petal fall and every 10 to 14 days thereafter until control is achieved.	
Peach, Apricot	codling moth	Start to spray when green leaf tips appear. Repeat at 10 day intervals until bloom, and at 14 day intervals after bloom.	
Nectarine, Quince, Persimmon, Pomegranate	Japanese Beetle, June beetle, orange tortrix, tarnished plant bug, tussock moth, red spider mite, aphid		
Cherries, Plums, Prunes	black cherry aphid, mealy plum aphid, cherry fruit fly, Japanese beetle, tent caterpillar, orange tortrix, tussock moth		
	codling moth, plum curculio		
Shade Trees	red spider mite, flea beetle, Japanese beetle, leaf roller, birch leaf miner, lace bugs, tent caterpillar	Start to spray when green tips appear. Repeat at 10 day intervals.	
Asparagus	aphids, asparagus beetle	Apply when insects appear. Repeat as necessary.	6 to 12 lb. in 100 gal. per acre.
Beets, Radishes, Carrots	flea beetle, aphids, carrot rust fly		
Broccoli, Brussels Sprouts, Cabbage, Cauliflower	imported cabbage worm, cabbage looper, diamond back moth, aphids		
Tomatoes, Potatoes, Eggplant, Peppers, Pimientos, Sweet potatoes	Colorado potato beetle, flea beetle, leafhopper, young tomato hornworm, tomato pinworm, tarnished plant bug, aphid, white fly, mealybugs	Apply when plants are 3 inches high. Repeat at 5 to 7 day intervals.	10 to 20 lb. in 100 gal. per acre
Beans	red spider mite, Mexican bean beetle, leafhopper, stink bugs, tarnished plant bug, aphids, bean leaf beetle.		

CROP	INSECTS	WHEN TO TREAT	AMOUNT
Celery	parsnip webworm, two-spotted spider mite, tarnished plant bug, carrot rust fly, aphid	Apply thoroughly to plants as soon as they appear above ground. Repeat at 3 to 7 day intervals.	5 to 10 lb. in 100 gal. per acre
Cucumber, Melon, Pumpkin, Squash (summer & winter), Watermelon, Cantaloupe	pickleworm, melonworm, cucumber beetle, young squash bug, aphids		
Lettuce, Collard, Kale, Mustard Greens, Spinach, Swiss Chard, Kohlrabi, Okra, Endive, Salsify	flea beetle, harlequin bug, leafhoppers, cabbage worm, looper, stink bug, white fly, lygus bug		
Peas, Blackeyed Peas, Cowpeas	aphids, leafhopper		
Turnips, Rutabagas, Parsnips, Horseradish	flea beetle, harlequin bug, leafhopper, lygus and stink bugs, cabbage worm, parsnip webworm		
Flowers, Shrubs	aphids, mites, rose chafer, Japanese beetle, flea beetle, leafhopper, red spider mite and repeat bagworm	Start to spray when insects appear. Repeat as necessary.	5 to 10 lb. in 100 gal. per acre
Blackberry, Raspberry, Dewberry, Loganberry, Boysenberry, Youngberry	raspberry sawfly, red spider leafminer	A week before blossoms open.	7 to 15 lb. in 100 gal. per acre
	raspberry fruit worm, cane borer	10 days after blossoming.	
Blueberry, Huckleberry, Grape	flea beetle, grape leaf skeletonizer, Japanese beetle, leaf hopper, rose chafer	When insects first appear. Repeat as necessary.	
	blueberry maggot	When most of fruit is set.	
Current, Gooseberry	Imported currant worm	When most of fruit is set. Repeat as necessary.	
	aphid, red spider mite	When insects first appear. Repeat as necessary.	
Strawberry	leaf roller, weevil, red spider mite	Before and after blossoming.	
	spittlebug	When frothy spittle masses appear. Repeat as needed.	

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.