

100

Keep out of reach of children.

This material is toxic. Avoid inhaling dust, vapor or mist from sprays. Avoid contact with skin or eyes. Avoid contamination of food and drink. Respiratory vapor and dust may cause irritation of skin and eyes. May be absorbed through the skin if in solution. In case of skin contact wash with plenty of water and remove for first aid. Wash eyes with water and get prompt medical attention.

1924

UNDER THE FEDERAL BUREAU OF INVESTIGATION

UNDER THE FEDERAL GOVERNMENT



MoorMaFume®

A Wettable Powder Insecticide

FOR CONTROL OF

- Hog Lice and Sarcoptic Mange
- Beef Cattle Lice, Ticks, Sarcoptic Mange and Scab Mites
- Sheep and Goat Lice, Ticks (keds), Sarcoptic Mange and Scab Mites (scabies)

Active Ingredient:

Lindane.....6%
Inert Ingredients.....94%

CAUTION

Keep out of reach of children.

Use only as recommended in instructions. Observe additional cautions on back panel.

This material is toxic. Avoid inhaling dust, vapor or mist from sprays. Avoid contact with skin or eyes. Avoid contamination of feed and foodstuffs. Harmful vapor and dust may cause irritation of skin and eyes. May be absorbed through the skin if in solution. In case of skin contact wash with plenty of soap and water. For eyes, flush with water and get prompt medical attention.

Do not use in dwellings.

USDA REG. NO. 1157-9

NET WT. 5 LBS.

ACCEPTED

10/7/71

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND
FOR PESTICIDE CONTROL REGISTRY
ED UNDER NO.

1157-9

INSTRUCTIONS

FOR SARCOPTIC MANGE OR LICE IN HOGS (SPRAY OR DIP).

For young animals (over 3 months of age)—Mix ½ lb. of MoorMaFume with 12 gallons of water and use as a spray or dip.

For mature animals—Mix 1 lb. of MoorMaFume with 12 gallons of water and use as a spray or dip.

The sprayer should be agitated or shaken occasionally during spraying to keep the MoorMaFume in suspension. Spray the hogs thoroughly, making sure all affected areas including the inside of the pig's ears and the underside of its body, are well soaked. One gallon of spray will treat 8 to 10 pigs weighing 100 lbs.

FOR HOG LICE (SPRINKLE).

Sprinkle evenly at the rate of 2 oz. to each 50 sq. ft. in the bedding and on all places where hogs sleep. Be sure all the hogs sleep in the treated areas. Allow ventilation but avoid drafts. For a thorough job repeat treatment in two weeks. Do not apply directly to animals.

FOR BEEF CATTLE, SHEEP AND GOATS—LICE, TICKS (KEDS), SCAB MITES (SCABIES), SARCOPTIC MANGE AND HORN FLIES (SPRAY OR DIP).

For young animals (over 3 months of age)—Mix ½ lb. of MoorMaFume with 12 gallons of water and use as a spray or dip.

For mature animals—Mix 1 lb. of MoorMaFume with 12 gallons of water and use as a spray or dip.

Agitate occasionally to maintain a good suspension. Allow one to two quarts per animal. One application is generally sufficient. MoorMaFume kills lice eggs as well as the lice themselves by contact. For sarcoptic mange it is only necessary to treat thoroughly the affected areas, but for lice, ticks and scab mites, the entire animal should be thoroughly sprayed or dipped. For horn fly control in beef cattle, the treatment may be repeated every 2 to 3 weeks if necessary.

CAUTION. Do not treat animals under three months of age. Do not treat animals within 30 days of slaughter. Do not dip animals within 60 days of slaughter. Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

Do not treat dairy animals or use in dairy barns. Do not use on poultry or in poultry houses.

Do not reuse container. Destroy when empty.

Manufactured for Moorman Mfg. Co., Quincy, Illinois 62301

