



Rid-Ezy®

Medicated

A medicated mineral feed for cattle containing ronnel for continuous feeding during the summer for the control of horn flies and cattle grubs. A 14-day treatment for grub control only.

For beef cattle and non-lactating dairy animals only. Follow feeding directions carefully. To be used as sole source of ronnel medication.


CAUTION: Keep out of reach of children.



WARNING: Will draw 10 days prior to slaughter.

Observe additional cautions and warnings on back panel.

U.S. Patent 2,599,516. Contents licensed for use under U.S. Patent 2,811,450
EPA Reg. No. 1157-24-AB



879

MOORMAN'S RID-EZY MEDICATED

Moorman Mfg. Co., General Office, Quincy, Illinois

A medicated mineral feed for cattle containing ronnel for continuous feeding during the summer for the control of horn flies and cattle grubs. A 14-day treatment for grub control only.

For beef cattle and non-lactating dairy animals only. Follow feeding directions carefully. To be used as sole source of ronnel medication.

Active ingredient: Ronnel (p, O-Dimethyl 0-(2, 4, 5-trichlorophenyl) Phosphorothioate) 8.5%
Inert ingredients 91.5%

GUARANTEED ANALYSIS

Calcium (Ca), not less than 13.0%
Calcium (Ca), not more than 18.6%
Phosphorus (P), not less than 8.0%
Salt (NaCl), not less than 27.0%
Salt (NaCl), not more than 32.0%
Iodine (I), not less than 0.016%

INGREDIENTS

Low Fluorine Ground Rock Phosphate, Dicalcium Phosphate, Dehydrated Phosphate, Limestone, Salt, Sulfate of Iron, Iron Oxide, Potassium Iodide, Calcined Diatomaceous Earth, Cobalt Carbonate, Copper Carbonate, Manganese Oxide, Magnesium Oxide, Zinc Oxide, Potassium, Cane Molasses, Soybean Meal.

Observe additional cautions and warnings on back panel.

U.S. Patent 2,599,516. Contents licensed for use under U.S. Patent 2,811,480
EPA Reg. No. 1157-24-AB

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
EPA NUMBER NO. 1157-24

879

**MOORMAN'S RID-EZY
MEDICATED**

Moorman Mfg. Co., General Office, Quincy, Illinois

A medicated mineral feed for cattle containing ronnel for continuous feeding during the summer for the control of horn flies and cattle grubs. A 14-day treatment for grub control only.

For beef cattle and non-lactating dairy animals only. Follow feeding directions carefully. To be used as sole source of ronnel medication.

Active Ingredient: Ronnel [O, O-Dimethyl O-(2, 4, 5-trichlorophenyl) Phosphorothioate] 5.5%
Inert Ingredients 94.5%

GUARANTEED ANALYSIS

Calcium (Ca), not less than 13.0%
Calcium (Ca), not more than 15.6%
Phosphorous (P), not less than 5.0%
Salt (NaCl), not less than 27.0%
Salt (NaCl), not more than 32.0%
Iodine (I), not less than 0.016%

INGREDIENTS

Low Fluorine Ground Rock Phosphate, Dicalcium Phosphate, Defluorinated Phosphate, Limestone, Salt, Sulfate of Iron, Iron Oxide, Potassium Iodide, Calcined Diatomaceous Earth, Cobalt Carbonate, Copper Carbonate, Manganous Oxide, Magnesium Oxide, Zinc Oxide, Petrolatum, Cane Molasses, Soybean Meal.

MoorMan's Rid-Ezy Medicated

Instructions for feeding

Observe warning and caution statements

Note: Horn flies may develop resistance to ronnel through sub-lethal exposure to various organic phosphates. Should horn fly control prove unsatisfactory when Rid-Ezy is used according to instructions, change to a fly control program involving an active ingredient other than ronnel.

Continuous feeding for horn fly and grub control:

When to feed—Start feeding in spring or summer before horn flies appear on the cattle and after heel fly activity has stopped. Continue feeding all summer and into the fall until cold weather restricts fly development. Feed continuously for a minimum of 75 days for grub control. Southern areas may require feeding earlier in the spring and continuing later into the fall than do northern areas.

How much is needed—Recommended consumption of Rid-Ezy is $\frac{1}{4}$ lb. per 100 lbs. of body weight per head per month.

How to feed—self-feeding—Self-feed continuously near all watering and loafing areas so that cattle have constant access to Rid-Ezy.

Provide one mineral feeder box for each 15 to 20 head so all cattle will have an equal opportunity to eat Rid-Ezy. Keep feeders, troughs, bunks clean and put out only a 5 to 7 day supply at one time to encourage even, normal consumption of Rid-Ezy. Protection of Rid-Ezy from heavy rains is desirable. Salt or other mineral is not recommended when cattle are being self-fed Rid-Ezy.

How to feed—mixed ration feeding—When cattle are on supplemental feeds, or when Rid-Ezy is not consumed at recommended levels when self-fed, Rid-Ezy should be mixed as directed with protein supplement, ground grain, or a mixture of both, and fed daily.

Under these conditions, mix Rid-Ezy in the supplemental feed to provide 0.008 lb. Rid-Ezy per hundred pounds body weight per day (0.25 lb. per hundred pounds body weight per head per month).

Example: 100 animals averaging 500 lbs. would require 4 lbs. Rid-Ezy daily mixed in the feed. (100 animals X 5 hundred-weight X 0.008 lb. Rid-Ezy = 4 lbs. Rid-Ezy per day.)

Do not mix more than a 7-day supply at one time.

Do not allow cattle to run out of Rid-Ezy. Feed continuously all summer or for at least 75 days for grub control.

When not to feed—Do not feed to sick animals. Do not feed to animals under stress, such as those just shipped or dehorned, castrated or weaned within the last 10 days.

14-day feeding treatment for grub control only:

Rid-Ezy may be mixed with feed (grain or protein supplement) and fed continuously for 14 days.

When to feed—Start feeding any time after heel fly activity has stopped but before grubs have started to cut in the back. Southern area—June thru September; Northern area—August thru December. Feed Rid-Ezy only when cattle are in good condition and eating well.

How much is needed—The 14-day treatment requires 0.2 lb. Rid-Ezy for each 100 lbs. cattle weight.

How to mix—Mix each 1 lb. of Rid-Ezy with 20 lbs. other feed to be called the "Mixture (Feed Plus Rid-Ezy)". Do not mix more feed than will be fed in 14 days.

How to feed—Feed daily 0.3 lb. of "Mixture (Feed Plus Rid-Ezy)" per 100 lbs. cattle weight for 14 continuous days.

Make certain that all cattle are eating and that adequate feeding space is available so each animal can get its proportionate share. Since cattle normally are being fed more feed daily than the recommended amount of the "Mixture (Feed Plus Rid-Ezy)," make up the difference with non-medicated feed. Always make certain that the medicated feed is evenly distributed in the total ration.

When not to feed—Do not feed to sick animals. Do not feed to animals under stress, such as those just shipped or dehorned, castrated or weaned within the last 10 days. Do not feed to animals in which the grubs have started to cut in the back.



WARNING: For beef cattle and non-lactating dairy animals only. Withdraw 10 days prior to slaughter. Withdraw from dairy animals 10 days before calving; if dairy cows or heifers freshen during medication, or if medication has not been withdrawn the required 10 days prior to freshening, milk must not be used for food for 10 days after the last treatment. Ronnel is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

CAUTION

Keep out of reach of children. Wash hands with soap and water after handling. This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.