



Rid-Ezy[®] Block Medicated

ACCEPTED
JUN 3 1974
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 1157-26

A medicated mineral feed for cattle containing ronnel for continuous feeding during the summer for the control of horn flies and cattle grubs.

For beef cattle and non-lactating dairy animals only. Follow feeding directions carefully. To be used as sole source of ronnel medication.

Active Ingredient:
Ronnel [O,O-Dimethyl-O-(2,4,5-trichlorophenyl) Phosphorothioate] 5.5%
Inert ingredients 94.5%

INGREDIENTS

Low fluorine ground rock phosphate	Salt	Copper carbonate	Soybean meal
Dicalcium phosphate	Sulfate of iron	Manganous oxide	Dextrinized starch
Defluorinated phosphate	Iron oxide	Magnesium oxide	Paraffin wax
Limestone	Potassium iodide	Zinc oxide	Hemicellulose extract
	Cobalt carbonate	Cane molasses	

GUARANTEED ANALYSIS

Calcium (Ca), not less than 13.0%	Phosphorus (P), not less than 5.0%	Salt (NaCl), not more than 32.0%
Calcium (Ca), not more than 15.6%	Salt (NaCl), not less than 27.0%	Iodine (I), not less than 0.016%



CAUTION—Keep out of reach of children.

WARNING: Withdraw 10 days prior to slaughter.

Observe additional cautions and warnings on side panel.

U. S. Patent 2,500,516

Contacts licensed for use under U. S. Patent 2,811,480 • EPA REG. No. 1157.26.AB



Rid-Ezy[®] Block Medicated

ACCEPTED

JUN 3 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 1157-26

A medicated mineral feed for cattle containing ronnel for continuous feeding during the summer for the control of horn flies and cattle grubs.

For beef cattle and non-lactating dairy animals only. Follow feeding directions carefully. To be used as sole source of ronnel medication.

Active ingredient:

Ronnel [O,O-Dimethyl-O-(2,4,5-trichlorophenyl) Phosphorothioate] 5.5%

Inert ingredients 94.5%

INGREDIENTS

Low fluorine ground rock
phosphate
Dicalcium phosphate
Defluorinated phosphate
Limestone

Salt
Sulfate of iron
Iron oxide
Potassium iodide
Cobalt carbonate

Copper carbonate
Manganous oxide
Magnesium oxide
Zinc oxide
Cane molasses

Soybean meal
Dextrinized starch
Paraffin wax
Hemicellulose extract

GUARANTEED ANALYSIS

Calcium (Ca), not less than 13.0%
Calcium (Ca), not more than 15.6%

Phosphorus (P), not less than 5.0%
Salt (NaCl), not less than 27.0%

Salt (NaCl), not more than 32.0%
Iodine (I), not less than 0.016%



CAUTION—Keep out of reach of children.

WARNING: Withdraw 10 days prior to slaughter.

Observe additional cautions and warnings on side panel.

U. S. Patent 2 500 516

Contents licensed for use under U. S. Patent 2 811 180

FPA REG. No. 1157-26-AR

EPH

11-40-7

MoorMan's Rid-Ezy Block Medicated

INSTRUCTIONS FOR FEEDING FOR HORN FLY AND GRUB CONTROL OBSERVE WARNING AND CAUTION STATEMENTS



Note: Horn flies may develop resistance to ronnel through sub-lethal exposure to various organic phosphates. Should horn fly control prove unsatisfactory when Rid-Ezy is used according to instructions, change to a fly control program involving an active ingredient other than ronnel.

When to feed—Start feeding in spring or summer before horn flies appear on the cattle and after heel fly activity has stopped. Continue feeding all summer and into the fall until cold weather restricts fly development. Feed continuously for a minimum of 75 days for grub control. Southern areas may require feeding earlier in the spring and continuing later into the fall than do northern areas.

How much is needed—Recommended consumption of Rid-Ezy is $\frac{1}{4}$ lb. per 100 lbs. of body weight per head per month.

How to feed—Self-feed continuously near all watering and loafing areas so cattle have constant access to Rid-Ezy. Put out one Rid-Ezy Block for each 15 cattle. Scatter the blocks so that all cattle have an equal opportunity to consume Rid-Ezy in recommended amounts.

Do not feed salt or other mineral when cattle are being fed Rid-Ezy.

Do not allow cattle to run out of Rid-Ezy. Feed continuously all summer or for at least 75 days for grub control.

When not to feed—Do not feed to sick animals. Do not feed to animals under stress, such as those just shipped or dehorned, castrated or bled within the last 10 days.



WARNING: For beef cattle and non-lactating dairy animals only. Withdraw 10 days prior to slaughter. Withdraw from dairy animals 10 days before calving; if dairy cows or heifers freshen during medication, or if medication has not been withdrawn the required 10 days prior to freshening, milk must not be used for food for 10 days after the last treatment. Ronnel is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

CAUTION—Keep out of reach of children. Wash hands with soap and water after handling.

This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

Store Rid-Ezy in a cool, dry place.