


NET WT. 12½ LBS.

MoorMan's

INSECTICIDE DUST

An aid in the control of horn flies and lice on beef cattle



ACTIVE INGREDIENTS: Methoxychlor Technical*3%
*Equivalent to 2.64% 2,2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane
and 0.36% other isomers and related compounds.
INERT INGREDIENTS.....97%

INSTRUCTIONS FOR USE

Self-application—Dust Bag

Place approximately 6 lbs. of Insecticide Dust (½ the contents of the paper bag container) in the burlap dust bag supplied with this container. Allow one (1) dust bag per 10 to 15 head of cattle. Hang these bags in a line or group them in areas which the cattle frequent, such as shade trees or gates and lanes used as pathways to watering or feeding areas. Bags should be hung at least 24 inches apart, and so that the bottom of the bag is about 30 inches from the ground. The recommended dosage is 12½ lbs. per 10 to 15 head of cattle per month. In case the recommended amount of dust is not used in a month's period, relocate bags to an area more accessible to the cattle. Do not use more than 1 lb. of Insecticide Dust per head per month.

Hand Application

Insecticide Dust may be applied to beef animals by sprinkling the Dust over the animals, or by using a suitable mechanical dust applicator. Animals should be thoroughly dusted once every week, if necessary, for horn fly and lice control. Do not use more than ¼ lb. of Dust per animal per treatment. Persons treating animals should avoid inhalation of Dust.

CAUTION

KEEP OUT OF REACH OF CHILDREN



Do not hang bags in areas accessible to children. Avoid inhalation of dust. Do not use on animals within 30 days of slaughter. Do not use on dairy animals. Do not use in or around dairy barns. Do not contaminate feed and water.

This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of waste.

U.S.D.A. REG. NO. 1157-29

MOORMAN MFG. COMPANY, GENERAL OFFICE, QUINCY, ILLINOIS

705