

NET WT. 12 1/2 LBS.

Moor-Man's

Dust with † Co-Ral Insecticide

For control of horn flies and lice on beef and dairy cattle, and lice on swine

ACTIVE INGREDIENT:

O, O-Diethyl O-(3-chloro-4-methyl-2-oxo(2H)-1-benzopyran-7-yl) Phosphorothioate 1%
INERT INGREDIENTS: 99%

INSTRUCTIONS FOR USE

Self-application—dust bag for beef and dairy cattle

Place approximately 6 lbs. of Dust with Co-Ral Insecticide (1/2 the contents of the paper bag container) in the burlap dust bag supplied with this container. Allow one (1) dust bag per 12 to 24 head of cattle. Hang these bags in a line or group them in areas which the cattle frequent, such as shade trees or gates and lanes used as pathways to watering or feeding areas. For lactating dairy cows, bags may be suspended only in the exit through which the cows leave the milking barn. Bags should be hung at least 24 inches apart so that the bottom of the bag will hang 4 to 6 inches below the topline of the cattle. The recommended dosage is 12 1/2 lbs. per 12 to 24 head of cattle per month.

In case the recommended amount of Dust is not used in a month's period, relocate bags to an area more accessible to the cattle. Do not use more than 1 lb. of Cattle Dust per head per month.

Direct application to swine for control of lice.

Apply up to 1 ounce per animal as a uniform coat to head, shoulders, and back. Repeat as necessary, but not more often than once every 10 days. No interval is required between treatment and use of meat as feed.

Hand application for beef and dairy cattle

Dust with Co-Ral Insecticide may be applied to beef and dairy cattle by sprinkling the Dust over the animals by use of a shaker can or by using a suitable mechanical dust applicator. Animals should be thoroughly dusted once every week, if necessary, for horn fly and lice control. No interval is required between treatment and use of meat or milk as food. Do not use more than 2 ounces (1/4 lb.) of Dust per animal per treatment. Persons treating animals should avoid inhalation of the dust and prolonged contact of the dust with the skin.

Direct application and bedding treatment for control of severe hog lice infestations.

Apply 2 ounces uniformly over each 30 square feet of fresh, dry bedding and not more than 1 ounce directly on each hog as a uniform coat to head, shoulders, and back. Repeat as necessary, but not more often than once every 10 days.

Restrictions:

Do not hang dust bags over or contaminate feed, mineral, water or milking equipment with Dust.

CAUTION: Keep out of reach of children

Harmful to humans if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe Dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate food, feed or water. Keep out of reach of children.

Destroy damaged or worn-out dust bag by burying or burning. Stay away from smoke or fumes.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression. To Physician: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered with atropine.

This product is toxic to fish, birds and other wildlife. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

†Co-Ral is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen
EPA REG. NO. 1157-31
EPA Est. 1157-IL-1

Moorman Mfg. Co., General Office, Quincy, Illinois 62301

Code 8

706

