



IGR MineralTM Block

INSTRUCTIONS FOR FEEDING

To prevent the breeding of horn flies in the manure of treated cattle. A pesticidally active mineral feed for beef cattle containing methoprene* insect growth regulator for continuous feeding during the fly season.

Make sure cattle are not starved for minerals or salt by providing them before feeding MoorMan's IGR Mineral Block. Then remove salt or mineral supplement and place MoorMan's IGR Mineral Block before cattle as the only source of minerals. Start feeding MoorMan's IGR Mineral Block continuously in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity.

| | |
|--|----------------|
| ACTIVE INGREDIENT: methoprene [Isopropyl (E,E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate]* | 0.02% |
| INERT INGREDIENTS: | 99.98% |
| TOTAL | 100.00% |

The recommended consumption of MoorMan's IGR Mineral Block is 0.25 lb. (¼ lb.) to 0.5 lb. (½ lb.) per 100 lbs. of bodyweight per month. Allow one MoorMan's IGR Mineral Block for each 15 head and place near watering or loafing areas. Protect blocks from rain.

GUARANTEED ANALYSIS

INGREDIENTS

Calcium (Ca), not less than 13.2%
 Calcium (Ca), not more than 15.8%
 Phosphorus (P), not less than 7.0%
 Salt (NaCl), not less than 25.0%
 Salt (NaCl), not more than 30.0%
 Iodine (I), not less than 0.015%
 Vitamin A, not less than 200,000 USP Units per pound.

Vitamin A Acetate, D-Activated Animal Sterol (Source of Vitamin D₃—Stability Improved), Dicalcium Phosphate, Defluorinated Phosphate, Calcium Carbonate, Low Fluorine Ground Rock Phosphate, Soybean Meal, Salt, Iron Sulfate, Iron Oxide, Potassium Iodide, Cobalt Carbonate, Silicon Dioxide, Copper Carbonate, Cane Molasses, Manganese Oxide, Ethylene Diamine Dihydrochloride, Zinc Oxide, Magnesium Oxide, Paraffin Wax, Hemicellulose Extract.

If consumption of MoorMan's IGR Mineral Block is above 0.5 lb. (½ lb.) per 100 lbs. of bodyweight per month, reduce the number of feeding locations where this block is fed or relocate blocks in areas less frequented by cattle.

If consumption of MoorMan's IGR Mineral Block is below 0.25 lb. (¼ lb.) per 100 lbs. of bodyweight per month, increase the number of feeding locations where this block is fed or relocate blocks in areas more frequented by cattle.

*As the insect growth regulator, ALTOSID, TM Reg. U.S. Pat. Office US Pats. 3,904,662 & 3,912,815

Use this product only as specified on the label.

CAUTION: Keep out of reach of children.



WARNING: Do not feed to beef cattle within 14 days of slaughter.



CAUTION: Keep out of reach of children.

WARNING: Do not feed to beef cattle within 14 days of slaughter.

TAKE TIME



OBSERVE LABEL DIRECTIONS

Moorman Mfg. Co., General Office, Quincy, Illinois 62301

EPA REG. NO. 1157-42 EPA Est. 1157-IL-1

Net. Wt. 50 Lbs.

CODE 1

ACCEPTED
 NOV 19 1975
 1157-42