

IGR MineralTM Block

To prevent the breeding of horn flies in the manure of treated cattle. A pesticidally active mineral feed for beef cattle containing methoprene* insect growth regulator for continuous feeding during the fly season.

ACTIVE INGREDIENT: methoprene [Isopropyl (E,E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate]* 0.02%
INERT INGREDIENTS: 99.98%
TOTAL 100.00%

GUARANTEED ANALYSIS

Calcium (Ca), not less than 13.2%
Calcium (Ca), not more than 15.8%
Phosphorus (P), not less than 7.0%
Salt (NaCl), not less than 25.0%
Salt (NaCl), not more than 30.0%
Iodine (I), not less than 0.015%
Vitamin A, not less than 200,000 USP Units per pound.

INGREDIENTS

Vitamin A Acetate, D-Activated Animal Sterol (Source of Vitamin D, -Stability Improved), Dicalcium Phosphate, Deffluorinated Phosphate, Calcium Carbonate, Low Fluorine Ground Rock Phosphate, Soybean Meal, Salt, Iron Sulfate, Iron Oxide, Potassium Iodide, Cobalt Carbonate, Silicon Dioxide, Copper Carbonate, Cane Molasses, Manganous Oxide, Ethylene Diamine Dihydriodide, Zinc Oxide, Magnesium Oxide, Paraffin Wax, Hemmcel lulose Extract.

*As the insect growth regulator, ALTOSID[®] a trademark of

ZOECON CORPORATION

US Pats. 3,904,662 & 3,912,815

Use this product only as specified on the label.

CAUTION: Keep out of reach of children.



WARNING: Do not feed to beef cattle within 14 days of slaughter.

Moorman Mfg. Co., General Office, Quincy, Illinois 62301

EPA REG. NO. 1157-42 EPA Est. 1157-IL-1

Net Wt. 50 Lbs.

INSTRUCTIONS FOR FEEDING

Make sure cattle are not starved for minerals or salt by providing them before feeding MoorMan's IGR Mineral Block. Then remove salt or mineral supplement and place MoorMan's IGR Mineral Block before cattle as the only source of minerals. Start feeding MoorMan's IGR Mineral Block continuously in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity.

The recommended consumption of MoorMan's IGR Mineral Block is 0.25 lb. (¼ lb.) to 0.5 lb. (½ lb.) per 100 lbs. of bodyweight per month. Allow one MoorMan's IGR Mineral Block for each 15 head and place near watering or loafing areas. Protect blocks from rain.

If consumption of MoorMan's IGR Mineral Block is above 0.5 lb. (½ lb.) per 100 lbs. of bodyweight per month, reduce the number of feeding locations where this block is fed or relocate blocks in areas less frequented by cattle.

If consumption of MoorMan's IGR Mineral Block is below 0.25 lb. (¼ lb.) per 100 lbs. of bodyweight per month, increase the number of feeding locations where this block is fed or relocate blocks in areas more frequented by cattle.

CAUTION: Keep out of reach of children.



WARNING: Do not feed to beef cattle within 14 days of slaughter.

TAKE TIME



OBSERVE LABEL DIRECTIONS

CODE 1

1157-42

1/5/1976

Page 122

2 7 2

B-50-B

MoorMan's

ACCEPTED

JAN 5 1976

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
can be used, the pesticide
registered under
EPA Reg. No. 1157-42

1167-42

650-B

MoorMan's