

# IGR·Mineral® Block

To prevent the breeding of horn flies in the manure of treated cattle. A pesticidally active mineral feed for beef and dairy cattle containing methoprene\* insect growth regulator for continuous feeding during the fly season.

**ACTIVE INGREDIENT:** methoprene [Isopropyl (E,E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate]\* ..... 0.02%  
**INERT INGREDIENTS:** ..... 99.98%  
**TOTAL** ..... 100.00%

## GUARANTEED ANALYSIS

Calcium (Ca), not less than ..... 13.2%  
Calcium (Ca), not more than ..... 15.8%  
Phosphorus (P), not less than ..... 7.0%  
Salt (NaCl), not less than ..... 25.0%  
Salt (NaCl), not more than ..... 30.0%  
Iodine (I), not less than ..... 0.015%  
Vitamin B<sub>12</sub>, not less than 200,000 USP Units per pound.

## INGREDIENTS

Vitamin A Acetate, D-Activated Animal Sterol (Source of Vitamin D<sub>3</sub>—Stability Improved), Dicalcium Phosphate, Defluorinated Phosphate, Calcium Carbonate, Low Fluorine Ground Rock Phosphate, Soybean Meal, Salt, Iron Sulfate, Iron Oxide, Potassium Iodide, Cobalt Carbonate, Silicon Dioxide, Copper Carbonate, Cane Molasses, Manganous Oxide, Ethylene Diamine Dihydriodide, Zinc Oxide, Magnesium Oxide, Paraffin Wax, Hemicellulose Extract.

\*As the insect growth regulator, ALTOSID®, a trademark of

**ZOECON CORPORATION**

US Pats. 3,904,662 & 3,912,815

Use this product only as specified on the label.

Keep out of reach of children.

**CAUTION**

Moorman Mfg. Co., General Office, Quincy, Illinois 62301

EPA REG. NO. 1157-42 EPA Est. 1157-IL-1

**Net Wt. 50 Lbs.**

## INSTRUCTIONS FOR FEEDING

Make sure cattle are not starved for minerals or salt by providing them before feeding MoorMan's IGR Mineral Block. Then remove salt or mineral supplement and place MoorMan's IGR Mineral Block before cattle as the only source of minerals. Start feeding MoorMan's IGR Mineral Block continuously in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity.

The recommended consumption of MoorMan's IGR Mineral Block is 0.25 lb. (¼ lb.) to 0.5 lb. (½ lb.) per 100 lbs. of bodyweight per month. Allow one MoorMan's IGR Mineral Block for each 15 head and place near watering or loafing areas. Protect blocks from rain.


If consumption of MoorMan's IGR Mineral Block is above 0.5 lb. (½ lb.) per 100 lbs. of bodyweight per month, reduce the number of feeding locations where this block is fed or relocate blocks in areas less frequented by cattle.

If consumption of MoorMan's IGR Mineral Block is below 0.25 lb. (¼ lb.) per 100 lbs. of bodyweight per month, increase the number of feeding locations where this block is fed or relocate blocks in areas more frequented by cattle.

**TAKE TIME**



**OBSERVE LABEL DIRECTIONS**



1157-42  
4/26/1977  
Page 182

