

1157-42

650-B



IGR Mineral® Block

To prevent the breeding of horn flies in the manure of treated cattle.
A pesticidally active mineral feed for beef cattle containing methoprene*
insect growth regulator for continuous feeding during the fly season.

ACTIVE INGREDIENT: methoprene [Isopropyl(E,E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate]* 0.02%
INERT INGREDIENTS: 99.98%
TOTAL 100.00%

GUARANTEED ANALYSIS

Calcium (Ca), not less than	13.2%
Calcium (Ca), not more than	15.8%
Phosphorus (P), not less than	7.0%
Salt (NaCl), not less than	25.0%
Salt (NaCl), not more than	30.0%
Iodine (I), not less than	0.015%
Vitamin A, not less than 200,000 International Units per pound.	

FEED INGREDIENTS

Vitamin A Acetate, D-Activated Animal Sterol (Source of Vitamin D₃—Stability Improved), Dicalcium Phosphate, Defluorinated Phosphate, Calcium Carbonate, Soybean Meal, Salt, Ferrous Sulfate, Iron Oxide, Calcium Iodate, Cobalt Carbonate, Silicon Dioxide, Copper Carbonate, Mineral Oil, Cane Molasses, Manganous Oxide, Ethylenediamine Dihydriodide, Zinc Oxide, Magnesium Oxide, Paraffin Wax, Hemicellulose Extract, Sodium Selenite.

*As the insect growth regulator, ALTOSID® a trademark of

ZOECON CORPORATION

US Pats. 3,804,662 & 3,812,815

Use this product only as specified on the label.

Keep out of reach of children.

CAUTION

Moorman Mfg. Co., General Office, Quincy, Illinois 62301-3488

EPA REG. NO. 1157-42 EPA Est. 1157-IL-1

650-B

MoorMan's

INSTRUCTIONS FOR FEEDING

Make sure cattle are not starved for minerals or salt by providing them before feeding MoorMan's IGR Mineral Block. Then remove salt or mineral supplement and place MoorMan's IGR Mineral Block before cattle as the only source of minerals. Start feeding MoorMan's IGR Mineral Block continuously in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity.

The recommended consumption of MoorMan's IGR Mineral Block is 0.25 lb ($\frac{1}{4}$ lb) to 0.5 lb ($\frac{1}{2}$ lb) per 100 lb of body weight per month. Allow one MoorMan's IGR Mineral Block for each 15 head and place near watering or loafing areas. Protect blocks from rain.

If consumption of MoorMan's IGR Mineral Block is above 0.5 lb ($\frac{1}{2}$ lb) per 100 lb of body weight per month, reduce the number of feeding locations where this block is fed or relocate blocks in areas less frequented by cattle.

If consumption of MoorMan's IGR Mineral Block is below 0.25 lb ($\frac{1}{4}$ lb) per 100 lb of body weight per month, increase the number of feeding locations where this block is fed or relocate blocks in areas more frequented by cattle.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal.

STORAGE: Store in dry area. Do not contaminate with pesticides or fertilizer.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty the plastic wrap, placing the block into application equipment. Then dispose of plastic wrap in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

ACCEPTED

AUG 28 1987

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 115-7-4/2