

inorians

IGR Mineral® Block

To prevent the breeding of horn files in the manure of treated cattle only.

A pesticidally active mineral feed for beef cattle containing methoprene* insect growth regulator for continuous feeding during the fly season.

ACTIVE INGREDIENT: methoprene [Isopropyl (E, E)-11-methoxy-3,7,11trimethyl-2, 4-dodecadienoate)* 0.02% INERT INGREDIENTS: 99.98%

HADANTEEN AMAI VOIC

GUARANI EED ANALTSIS	
Calcium (Ca), not less than 13.29	6
Calcium (Ca), not more than 15.89	
Phosphorus (P), not less than 7.01	Ž
Salt (NaCl), not less than 25.09	Ž
Salt (NaCl), not more than 30.01	
lodine (i), not less than 0.0159	
Vitamin A not lose than 200 00	š
Vitamin A, not less than 200,00	U

FEED INGREDIENTS

TOTAL 100.00%

Vitamin A Acetate, D-Activated Animal Steroi (Source of Vitamin D₃), Monocalcium Phosphate, Dicalcium Phosphate, Defluorinated Phosphate, Calcium Carbonate, Soybean Meal, Salt, Ferrous Sulfate, Iron Oxide, Potassium lodide, Cobalt Carbonate, Silicon Dioxide, Copper Sulfate, Mineral Oil, Cane Molasses, Manganous Oxide, Etylenediamine Dihydriodide, Zinc Oxide, Magnesium Oxide, Paraffin Wax, Hemicellulose Extract, Sodium Selenite.

"As the most growth regulator, ALTOSID," a trademark of Sandoz Ltd.

Use this product only as specified on the label. Keep out of reach of children.

CAUTION

Net Wt. 50 lb

Moorman Manufacturing Company, General Office Quincy, Illinois 62305-3115 EPA REG. NO. 1157-42 EPA Est. 1157-IL-1

MoorMan **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Make sure cattle are not starved for minerals or salt by providing them before feeding MoorMan's IGR Mineral Block. Then remove salt or mineral supplement and place MoorMan's IGR Mineral Block before cattle as the only source of minerals. Start feeding MoorMan's IGR Mineral Block continuously in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity.

The recommended consumption of MoorMan's IGR Mineral Block is 0.25 to (1/4 to) to 0.5 lb (1/2 lb) per 100 lb of body weight per month. Allow one MoorMan's IGR Mineral Block for each 15 head and place near watering or loafing areas. Protect blocks

If consumption of MoorMan's IGR Mineral Block is above 0.5 to (1/2 tb) per 100 to of body weight per month, reduce the number of feeding locations where this block is fed or relocate blocks in areas less frequented by cattle.

If consumption of MoorMan's IGR Mineral Block is below 0.25 lb (1/4 lb) per 100 lb of body weight per month, increase the number of feeding locations where this block is fed or relocate blocks in areas more frequented by cattle.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal.

STORAGE: Store in dry area. Do not contaminate with pesticides or fertilizer.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved yaste disposal facility...*

CONTAINER DISPOSAL: Completily ellipty the plastic weap, placing the block into application equipment. Then dispose of plastic wase in a sanitary landfill or by incineration, or, if allowed by State and Boas authorities, by Curning. If burned, stay out of smoke.

PRODUCT NO. 650-B

Code 9

