

Front Panel



IGR Cattle

Mix

BEST DOCUMENT AVAILABLE

1157-45

To prevent the breeding of horn flies in the manure of treated cattle
A pesticidally active feed for cattle containing methoprene* insect growth regulator for continuous feeding during the fly season

ACTIVE INGREDIENT: methoprene (Isopropyl (E,E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate)*	0.02%
INERT INGREDIENTS:	99.98%
TOTAL	100.00%

GUARANTEED ANALYSIS		FEED INGREDIENTS	
Crude Protein, not less than	40.0%	Soybean Meal	Corn Distillers Dried Solubles
Crude Fat, not less than	1.5%	Molasses	Corn Gluten Feed
Crude Fiber, not more than	8.0%	Mineral Oil	Cane

*As the insect growth regulator ALTOGID® a trademark of

ZOECON CORPORATION
US Pats. 3,354,562 & 3,912,815

Use this product only as specified on the back of the bag
Keep out of reach of children.

CAUTION

Moorman Mfg. Co., Cordele Division, Cordele, Georgia 31015-1137
EPA REG NO 1157-45 EPA Est 1157 GA 1

LOT NO.

NET WT. 50 LB.

Back Panel

MoorMan's IGR Cattle Mix INSTRUCTIONS FOR FEEDING

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product only prevents emergence of adult horn flies from the manure of the treated cattle. Begin use in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity. Introduction of the product when significant numbers of adult flies are present requires the use of adulticide sprays or dusts.

The dosage of MoorMan's IGR Cattle Mix is proportional to body weight. The recommended consumption of this product is 0.25 lb (1/4 lb) to 0.5 lb (1/2 lb) per 100 lb of body weight per month (30 days). Feed by either of the following methods:

Top-dress and hand-mix feeding

Top dress or hand-mix with supplemental feeds to provide 0.25 lb (1/4 lb) to 0.5 lb (1/2 lb) of IGR Cattle Mix per 100 lb body weight per month. When top dress feeding IGR Cattle Mix, have sufficient bunk space to allow all animals to eat at the same time.

Use the following table as a guide for determining the daily feeding rate of IGR Cattle Mix.

Body Weight (pounds)	Amount of IGR Cattle Mix required per head per day	
	By Weight	By Volume
500 - 599	1.2 ounces (0.075 lb)	1/4 cup*
700 - 999	1.6 ounces (0.10 lb)	1/3 cup*
1000 - 1600	2.4 ounces (0.15 lb)	1/2 cup*

*Standard household measuring cup

Mixed-ration feeding

Thoroughly mix IGR Cattle Mix with the daily grain or supplement to provide 0.25 lb (1/4 lb) to 0.5 lb (1/2 lb) of IGR Cattle Mix per 100 lb of body weight per month.

Use the following table as a guide for determining the proper mixing rate.

Body Weight (pounds)	Daily Grain or Supplemental Feed Consumption (pounds)			
	2 - 4	5 - 10	11 - 22	23 - 46
	Pounds of IGR Cattle Mix Needed per Ton of Grain or Supplemental Feed			
500	41.5	16.6	•	•
600	50.0	20.0	•	•
700	58.3	23.3	•	•
800	66.6	26.6	•	•
900	75.0	30.0	13.6	•
1000	83.3	33.3	15.2	•
1100	91.6	36.6	16.6	7.9
1200	100.0	40.0	18.2	8.7
1300	108.3	43.3	19.7	9.4
1400	116.6	46.6	21.2	10.1
1500	125.0	50.0	22.7	10.9
1600	133.3	53.3	24.2	11.6

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal.

STORAGE: Store in dry area.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedure.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

BEST DOCUMENT AVAILABLE

**LOT
NO**