

TWIN LIGHT

NET WEIGHT ONE POUND

ACTIVE INGREDIENTS:

Rotenone75%
Other Cube Extractives	1.50%
Copper as metallic (derived from basic copper sulfate)	5.00%
INERT INGREDIENTS	92.75%

ROTENONE COPPER DUST CAUTION

USDA Reg. No. 1459-39

KEEP OUT OF REACH OF CHILDREN
SEE REAR PANEL FOR OTHER CAUTIONS
FOR MANY

Insects and Diseases of Vegetables

SEACOAST LABORATORIES, INC.
257 HIGHWAY 18 • EAST BRUNSWICK, N. J.

ACCEPTED
527
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RO...
FOR ECONOMIC...
ED UNDER NO...
TO ATTACHED COMMENTS.

DIRECTIONS

CROP	PESTS
Cucumbers Melons Squash	Striped and Spotted Cucumber Beetles, Downy Mildew, Anthracnose, Leaf Spot and Bacterial Blight.
String Beans Lima Beans	Mexican Bean Beetles, Downy Mildew and Anthracnose.
Cabbage Broccoli Cauliflower	Cabbage Loopers, Flea Beetles, Diamond Back Moth and Downy Mildew.
Eggplant	Aphids, Flea Beetles, and Fruit Rot.
Peppers Tomatoes	Young Tomato Hornworm.

Twin Light Rotenone-Copper Dust may be applied any time of the day, on either dry or moist foliage. Apply dust at 7 day intervals, beginning soon after plants are above ground. Dosages will vary depending on cultural practices, size of plants and dusting equipment. Amount of dust used will accordingly range from 1/2 pound to 1 pound per 1000 square feet or 20 to 40 pounds to the acre per application. For best results keep foliage covered. **CAUTION** Avoid breathing dust.

Note: Make last application no closer than 1 day before harvest on edible crops.