

RECORDED
1972
11-24-66

TWIN® LIGHT RAT-AWAY

WARFARIN KILLER

FOR

RATS and MICE

ACTIVE INGREDIENT:

Warfarin 3-(alpha-acetonylbenzyl) -	
4-hydroxycoumarin	.5%
INERT INGREDIENTS	99.5%
Total	100.0%

Net Contents 4 oz.

EPA Reg.
No. 1159-66

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE FOR OTHER CAUTIONS

MANUFACTURED FOR
SEACOAST LABORATORIES, INC.
EAST BRUNSWICK, N. J.

DIRECTIONS (Follow Carefully)

RATS: Mix thoroughly one part of Rat-Away Powder with 19 parts of a suitable bait material which is available, such as corn meal, rolled oats, dog food, laying mash, stock food, etc. A small amount of a bland oil, such as mineral, corn, peanut, or any other edible oil, may be used, up to 5%, when the bait is so dry that it might have a tendency to dust.

When perishable baits, such as ground meat, fish, vegetables or similar material are used, bait placements should be made in late afternoon and replaced daily with fresh bait.

Dogs and cats will actively search out baits containing meat and fish. Therefore baits must be placed so as to prevent feeding by these animals.

Baits should be placed in amounts of from 4 oz. up to 1 lb. in locations frequented by rats or where rats have been known to feed. Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes.

Baits should be examined regularly and replaced when completely eaten. If bait moulds, decomposes or becomes sour it should be replaced.

Baiting should continue as long as evidence of feeding is observed. Except in extremely heavy infestations this should be for at least 10 days. Where a continuous source of infestation is present from nearby dumps or fields, permanent bait stations should be used and bait placed in them as often as necessary.

DIRECTIONS CONT'D (Follow Carefully)

MICE: For controlling mice follow the same general directions as for rats except that the amount of bait placements should be smaller. Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8-12 foot intervals.

Mice are more difficult to control than rats and complete control may take a longer period of baiting for at least 15 uninterrupted days.

IMPORTANT

Baiting should continue as long as evidence of feeding is observed. Except in extremely heavy infestations this should be from five days to two weeks. Where a continuous source of infestation is present from nearby dumps or fields, permanent bait stations should be used and bait placed in them as often as necessary.

CAUTION

Rat-Away Powder contains as its active ingredient an anti-coagulant chemical therefore baits prepared with Rat-Away Powder must be protected from or placed where humans, livestock, poultry or pets will not eat them. In case baits are eaten, give a teaspoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately. "Do not reuse container. Destroy when empty."

NOTE FOR PHYSICIANS: When a human has been known to have ingested any Rat-Away Powder, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol.

Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label."