




**TWIN LIGHT**®

**GRANULAR  
CLORO**



**CONTAINS  
5% TECHNICAL CHLORDANE**

# CONTAINS 5% TECHNICAL CHLORDANE

for control of  
**JAPANESE BEETLE, ORIENTAL BEETLE,  
ASIATIC BEETLE LARVAE, WHITE GRUBS  
SOD WEBWORMS AND ANTS**

#### APPROXIMATE SPREADER SETTINGS

Pounds per 1000 Sq. Ft.	1	2 1/2
Cyclone .....	2	2 1/2
Scotts .....	2 1/2	3 1/2
Lawn Beauty .....	2 1/2	3 1/2

Win Light GRANULAR CLORO is a product specifically formulated for soil or lawn application by a fertilizer - lime spreader.

#### DIRECTIONS

**LAWNS:** to control Japanese Beetle, Oriental Beetle, Asiatic Beetle Larvae and White Grubs, use 5 pounds per 1000 square feet.

**Sod Webworms** — Use 1 to 2 1/2 lbs. per 1000 square feet, **Ants, Chiggers, Earwigs, Mole Crickets, Spiders, Ticks, Cutworms, Centipedes, Grasshoppers and Army worms**—Use 1 lb. per 1000 square feet.

Wetting down the lawn area following a treatment is suggested to obtain better penetration of the toxicant into the soil.

Note: Keep children and pets off treated areas until treated areas have been watered and have dried completely.

Apply Contents of This Bag to 4000  
SQUARE FEET For GRUBPROOFING up  
to 20,000 SQUARE FEET for SURFACE PESTS

EPA Reg. No. 1159-84

NET WEIGHT 20 POUNDS

#### ACTIVE INGREDIENT:

Technical Chlordane\* ..... 5.00%

INERT INGREDIENTS: ..... 95.00%

\*Equivalent to 3% Octachloro-4, 7-Methanetetrahydroindane and 2% related compounds on a weight basis.

## CAUTION

KEEP OUT OF REACH OF CHILDREN AND  
DOMESTIC ANIMALS.

Avoid inhalation.

Avoid contact with skin, eyes and clothing. In case of skin contact, wash with soap and water; For eyes, get medical attention.

Avoid contamination of feed and foodstuffs.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

#### FISH and WILDLIFE CAUTION

Toxic to fish and wildlife. Birds feeding on treated lawn may be killed. Do not contaminate any body of water nor apply when conditions favor runoff from area treated.

ACCEPTED

OCT 1984

**Ticks, Cutworms, Centipedes, Grasshoppers and Army worms—Use 1 lb. per 1000 square feet.**

Wetting down the lawn area following a treatment is suggested to obtain better penetration of the toxicant into the soil.

Note: Keep children and pets off treated areas until treated areas have been watered and have dried completely.

**Apply Contents of This Bag to 4000  
SQUARE FEET For GRUBPROOFING up  
to 20,000 SQUARE FEET for SURFACE PESTS**

EPA Reg. No. 1159-84

**NET WEIGHT 20 POUNDS**

**Avoid inhalation.**

**Avoid contact with skin, eyes and clothing. In case of skin contact, wash with soap and water; For eyes, get medical attention.**


**Avoid contamination of feed and foodstuffs.**

**Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.**

**FISH and WILDLIFE CAUTION**

**Toxic to fish and wildlife. Birds feeding on treated lawn may be killed. Do not contaminate any body of water nor apply when conditions favor runoff from area treated.**

ACCEPTED  
001 11-11-84  
UNITED STATES  
ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460  
EPA REG. NO. 1159-84



**SEACOAST LABORATORIES, INC**  
**257 HIGHWAY 18 - EAST BRUNSWICK, N. J. 08816**