



TWIN LIGHT 5% METHOXYCHLOR DUST



DIRECTIONS

CUCURBITS (Cucumbers, Squash, Melons, etc.):

For Cucumber Beetles use 20 to 40 pounds per acre. Begin dusting when plants are very young. For Squash vine borer, begin dusting about July 1, apply 4 times at weekly intervals.

BEANS: For, Mexican Bean Beetles, Leaf Hoppers and Corn Earworms apply when injury is first noticed. Use 20 to 40 pounds per acre. Repeat when necessary.

POTATOES and TOMATOES: For Colorado Potato Beetles and Flea Beetles use 30 to 40 pounds per acre, repeat applications as needed.

ACTIVE INGREDIENTS:

Methoxychlor Technical 5.00%

INERT INGREDIENTS:

Equivalent to 1.40% 2,2-bis (p-Methoxyphenyl) 1,1,1-Trichloroethane and 0.60% of other isomers and reaction products. 95.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed or inhaled.

May cause irritation of nose, throat and skin.

Avoid breathing dust.

Avoid contact with eyes, skin and clothing. In case of contact, flush with plenty of water.

For eyes, get medical attention.

Do not apply later than 7 days prior to harvest. Wash treated produce thoroughly.

NET WEIGHT 50 POUNDS

Manufactured By

SEACOAST LABORATORIES, INC.

~~100 PENNY STREET, NEW YORK 14, N.Y.~~
EAST BRUNSWICK, N.J.

Handwritten notes and a date stamp: 12-15-62