

dried.

SOIL APPLICATION: For Cutworms, Farwigs, Millipedes, Leaf Eating Ants, Sowbugs and Mole Crickets, dust the ground around Tomatoes, Beans, Peas (only at time of planting or transplanting) and Flowers, use 1 pound per 500 square feet. Repear as necessary at 10 day intervals.

For control of Japanese Beetle Grubs, Root Maggots, Wireworms and White Grubs on Cabbage, Onions and Radishes, use 1 pound per 1000 square feet and work the product into top 2 to 3 inches of soil when the sced bed is being prepared.

water; for eyes, get medical attention.

Avoid contamination of feed and foodstuffs.


Use only according to the directions on this label.

#### FISH and WILDLIFE CAUTION

Toxic to fish and wildlife. Birds feeding over treated lawn may be **Killed**. Do not contaminate any body of water nor apply when conditions favor runoff from area treated.

## SPREADER SETTINGS

ACCEPTED	SIREADER SEITHOS		
	SURFACE PESTS		GRUB PROOFING
	Pounds Per 1000 Sq. Ft.	1 <b>lb</b> .	2 lbs.
	Scotts -	<del></del>	5
FEB 25 to 1	Lawn Beauty - 18"	4	5 1/4
TIN. A SHORE OF CT	Jackson - 20"	22	30
LON DO UNOME POISON RESERVED.	Buch =2 - 20"	1/2	3/4
10 Unil 112/154-113	Cyclone	2.8	3.4



SEACOAST LABORATORIES, INC 257 HIGHWAY 18 - EAST BRUNSWICK, N.J. 08816

# TWIN LIGHT PROFESSIONAL CHINCH BUG KILLER

with ASPON\*

USDA reg. No. 1159-x

#### Controls LAWN CHINCH BUGS

Sod Webworms and Ants

Covers 10,000 square feet

#### ACTIVE INGREDIENTS:

0,0,0-Tetrapropyl dithiopyrpphosphate......7.00%

### INERT INGREDIENTS

\*\*Equivalent to 1.2% Octachloro-4,7-Methanotetrahydroindane and .8% related compounds on a weight basis.

•ASPON is Stauffer Chemical Company's Trade Mark for an insecticide.

#### DIRECTIONS

LAWNS and ORNAMENTAL TURF-Apply 25 pounds(contents of bag) of this product on 10,000 square feet(100'x 100") with a fertilizer spreader for control of CHINCH BUGS, SOD WEBWORMS, EARWIGS, TICKS, and CHIGGERS. Immediately after application, drench with water or sprinkle thoroughly.

NOTE: Do not allow persons or pets on treated lawn until completely dry. Birds feeding over treated lawn may be harmed.

#### **WARNING:**

KEEP OUT OF REACH OF CHILDREN
MAY BE FATAL IF SWALLOWED
AVOID BREATHING DUST
DO NOT GET IN EYES, ON SKIN, ON CLOTHING
IN CASE OF CONTACT, IMMEDIATELY FLUSH EYES WITH PLENTY OF
WATER, FOR SKIN WASH WITH SOAP AND WATER
AVOID CONTAMINATION OF FOODSTUFIS, DRINKING WATER, FISH PONDS, etc.

OBSERVE THE FOLLOWING PRECAUTIONS:
Avoid drift to adjoining food and forage crops. To protect
fish and wildlife, do not contaminate streams, lakes or ponds
with this material.
Clean up spills on walks or patios.
DESTROY EMPTY CONTAINER-INCINERATE-STAY OUT OF SMOKE

Approximate Spreader Setting Cyclone......2% Scotts.......3%

Lawn Beauty .... 34

NET WEIGHT 25 POUNDS

Manufactured By

SEACUAST LABORATORIES, IN ..

. / Highway 18, East Brunswick, No.