

(GRANULAR CLAY BASE)

CD

(1)

M

EPA Reg No 1159-117

FPA Est 1159 N 1 1

FOR CONTROL of

JAPANESE BEETLE GRUBS in

FEDERAL and STATE QUARANTINE PROGRAMS ONLY

#### DIRECTIONS

This product is formulated for soil or TURF application by a seed spreader.

Use at the rate of 25 POUNDS PER 10,000 SQUARE FEET FOR GRUBPROOFING 2½ pounds per 1,000 square feet

NOTE: Following treatment, sprinkle the TURF thoroughly with water to wash insecticide into turf Do not allow children or pets on treated TURF until the insecticide is washed off the grass into the soil and grass is completely divid

#### APPROXIMATE SPREADER SETTINGS

Pounds per 1001 1 30

Cyclone Seculto

ACTIVE INGREDIENT.

Technical Chlordane: ...... 10.00% INERT INGREDIENTS: ..... 90.00%

\*Equivalent to 6% Octachloro—4, 7—Me thanotetrahydroindane and 4% related compounds on a weight basis

# CAUTION

## KEEP OUT OF REACH OF CHILDREN

**♥**oid inhalation.

Atold contact with skin, eyes and clothing in case of skin contact, wash with soap and water. For eyes, get medical attention

五二本 food contamination of fixed and foodstaffs

only according to the directions on this place

FISH and WILDLIFE CAUTION

## (GRANULAR CLAY BASE)

EPA Reg. No. 1159 117

> EFA Est 1159 N.J. 1

FOR CONTROL of

JAPANESE BEETLE GRUBS in

## FEDERAL and STATE QUARANTINE PROGRAMS ONLY

#### DIRECTIONS

This product is formulated for soil or TURF application by a seed spreader.

Use at the rate of 25 POUNDS PER 10,000 SQUARE FEET FOR GRUBPROOFING

2½ pounds per 1,000 square feet

NOTE: Following treatment, sprinkle the TURF thoroughly with water to wash insecticide into turf. Do not allow children or pets on treated TURF until the insecticide is washed off the grass into the soil and grass is completely dried.

#### APPROXIMATE SPREADER SETTINGS

Pounds per 1000 Sq. Ft	2 ½ lbs.
Cyclone	2 75
Scotts	3.5
Lawn Beauty	3 7 5

The age, condition of spreader, speed of operation and roughness of TURF will cause variations in rates applied. These suggested settings are for your guidance only

NET WEIGHT 25 POUNDS

ACTIVE INGREDIENT.

\*Equivalent to 6% Octachloro—4, 7—Me thanotetrahydroindane and 4% related compounds on a weight basis

# CAUTION

## KEEP OUT OF REACH OF CHILDREN

loid inhalation.

void contact with skin, eyes and clothing. In case of skin contact, wash with soap and water; For eyes, get medical attention

A oid contamination of feed and foodstuffs

e only according to the directions on this laber

#### FISH and WILDLIFE CAUTION

प्रिंड है । kxic to fish and wildlife. Birds feeding over treated areas may be killed. Do not contain inate any body of water not apply when conditions favor runoff from area treated.

Do not contaminate any body of water by direct application, cleaning of equipment, or disposar of wastes. Do not reuse empty bag. Destroy it by burying with waste or burning Stay away from smoke or fumes

Manufactured By

田

H

A

田

K

1976

# SEACOAST LABORATORIES, Inc.

EAST BRUNSWICK, N. J. 08816

NOTE: Following thoroughly with turf. Do not allow TURF until the institute the soil and APPROXII