

TWIN LIGHT

USDA Reg. No. 1159-120

CRABGRASS KILLER

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE REAR FOR ADDITIONAL CAUTIONS



Twin Light Liquid Crabgrass Killer will control Crabgrass in either the young or mature stages. The early treatment is preferable because the seedlings are easier to kill and the early control of crabgrass provides healthier grass in the late growing season.

DIRECTIONS FOR LAWNS AND FAIRWAYS

FOR YOUNG CRABGRASS: Apply with a garden hose jar sprayer (which attaches to garden hose) or a Pressure Sprayer during late Spring or early Summer. Two applications at 5 to 7 day intervals are generally needed for control. Repeated applications may be necessary if new crabgrass growth appears later in the season. Dilute with water and apply to measured area according to following table.
Liquid Crabgrass Killer

Required	Spray in gals. water	Area of Lawn Sq. Ft.
6 oz.	3.75	1000 Sq. Ft.
1 Pint	10	2000 Sq. Ft.
2 Gals.	50 to 100	1 Acre

FOR CRABGRASS IN THE BRANCH STAGE:

~~FOR MATURE CRABGRASS:~~ Apply with Garden Hose Spray jar (which attaches to garden hose) or pressure sprayer. Spray thoroughly to wet all the Mature Crabgrass plants. Two applications at 5 to 7 day intervals are generally required for control. Dilute with water and apply to measured area according to the following table.
Liquid Crabgrass Killer

Required	Spray in gals. water	Area of Lawn Sq. Ft.
8 oz.	5	1000 Sq. Ft.
1 Pint	10	2000 Sq. Ft.
2 1/2 Gals.	50 to 100	1 Acre

TEMPERATURE: If temperature is above 80°F., follow the dilution rates shown for young crabgrass (see above chart).

NOTE: Twin Light Liquid Crabgrass Killer is selective at the recommended rates of application in the presence of Kentucky, Merion Blue, Zoysia and Bermuda grasses. Temporary discoloration may occur in Fescue and Bent grass areas. Do not use on St. Augustine or Centerpeded grasses as injury may result. Above dosage also controls common chickweed, yellow foxtail, Goose Grass and Dallis Grass.

NOTE: IMPORTANT INFORMATION

- 1) Be sure soil is moist before treatment. During dry spells water thoroughly before applications are made.
- 2) Do not water lawn immediately after application. Do not apply if rain is expected within 24 hours after application.
- 3) Newly seeded lawns should not be treated until the third mowing.

CAUTION: Harmful if swallowed

CAUTION: Do not get in eyes or on the skin. If spilled on skin, wash off immediately with warm water and soap. Flush eyes with clear water. Avoid inhalation of spray mist. Avoid contamination of feed, foodstuffs, streams or other water sources. Keep beyond the reach of children and domestic animals. Keep livestock off treated areas. Wash hands after using.

ANTIDOTE: If taken internally, induce vomiting and call a doctor at once.

Licensed for use in the control of crabgrass according to the method of Patent 2,678,265 by reason of the inclusion in the purchase price of a royalty. A license at the same royalty rate to use the patented method with material obtained from any source will be granted, upon request, by Trustees of the U. S. Patent 2,678,265, 382 Springfield Avenue, Summit, N. J.


SEACOAST LABORATORIES, INC.

257 HIGHWAY 18 • EAST BRUNSWICK, N. J. 08816

ACCEPTED

MAY 2 1969

1156-120



Directions for use: Keep out of reach of children. Do not use on lawns or fairways. The only treatment for crabgrass is to prevent it from growing. Use this product on lawns and fairways to prevent crabgrass from growing.

DIRECTIONS FOR LAWNS AND FAIRWAYS

Apply this product to lawns and fairways in the spring and fall. It will prevent crabgrass from growing. Do not use on lawns or fairways that are already growing.

3.15

FOR CRABGRASS IN THE BRANCH STAGE

IMPORTANT INFORMATION: This product is a selective herbicide. It will kill crabgrass but not other grasses. Do not use on lawns or fairways that are already growing.

IMPORTANT INFORMATION

CAUTION: Harmful if swallowed

SEACOAST LABORATORIES, INC.

257 HIGHWAY 18 • EAST BRUNSWICK, N.J. 08817