

1159-122

ACC
APR 12, 1971
INDEXED
FUNGICIDE AND
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 1159109
TO ATTACHED DOCUMENTS.

Twin Light ROSE FUNGICIDE is recommended for the control of Black Spot and POWDERY MILDEW of ROSES and LEAF SPOT on IRIS and CHRYSANTHEMUMS.


DIRECTIONS

DILUTION: Use 1 1/2 level tablespoonsful per gallon of water (1 1/4 pounds per 100 gallons of water). May be used in a pressure sprayer or a bottle sprayer attached to a hose. Mix the required amount of Twin Light Rose Fungicide in a little water to make a thin paste, then add the necessary amount of water. Agitate during application.

<u>ROSES</u>	Black Spot Powdery Mildew	Make first application when young leaves first appear. Repeat at 7 to 10 day intervals, or as necessary throughout the season.
<u>IRIS</u> <u>CHRYSAN-</u> <u>THEMUMS</u>	Leaf Spot	Start treatments when plants are 6 inches high and repeat at 7 to 10 day intervals, or as necessary throughout the season.

NOTE: For COMBINATION INSECT and DISEASE CONTROL add 2 teaspoonsful of Twin Light Garden Spray to the diluted ROSE FUNGICIDE Spray.

COMPATIBILITY: Twin Light ROSE FUNGICIDE may be combined with the most commonly used insecticides and fungicides with the exception of strongly alkaline materials such as hydrated lime.



ACTIVE INGREDIENT: By Wgt.

Folpet *N (trichloromethylthio) phthalimide 75%

INERT INGREDIENTS: 25%

*PHALTAN is a registered trademark for phthalimide fungicide.

CAUTION

KEEP OUT OF REACH OF CHILDREN
Avoid inhalation of dust or spray mist. Avoid prolonged or repeated contact with skin. Wash hands after use. Avoid contamination of feed and foodstuffs. Twin Light ROSE FUNGICIDE should not be used in combination with or closely following oil sprays. Dispose of container promptly when empty. Do not reuse.

NOTE: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Manufactured for
SEACOAST LABORATORIES, INC.
EAST BRUNSWICK, N. J.

