

ACCEPTED

JUL 21 1960

PRECAUTIONARY STATEMENTS

**Hazards to Humans
(and Domestic Animals)**

CAUTION

AVOID BREATHING OF DUST, AVOID CONTACT WITH SKIN, AND CLOTHING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. WASH THOROUGHLY AFTER HANDLING. HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL THE FOLLOWING AFTERNOON.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Keep out of streams, rivers, or ponds. Do not contaminate water by use of equipment or disposal of wastes. Apply this product as specified on this label.

This pesticide is highly toxic to bees exposed to direct contact or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or blooming in the treatment area. Applications should be made to provide the maximum possible interval between applications and the next period of bee activity.

Directions for Use:

Violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE—Keep out of reach of children, pets and other animals. Avoid moisture. Avoid contamination of feed or foodstuffs by storage or disposal. Open burning and incineration prohibited. Do not reuse empty container.

DISPOSAL—Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State, or local disposal authorities.

CONTAINER DISPOSAL—Do not reuse empty container or container and put in trash collection.

NOT FOR USE IN THE HOME



***DIAZINON®
2% DUST**

**For Control of Certain INSECTS
That Infest VEGETABLES**

ACTIVE INGREDIENTS: O, O-diethyl O- (2-isopropyl 6-methyl-4-pyrimidinyl) phosphorothioate 2.00%
INERT INGREDIENTS 98.00%

*Diazinon® Trademark of CIBA-GEIGY Corp

**KEEP OUT OF REACH OF CHILDREN
CAUTION
STATEMENT OF
PRACTICAL TREATMENT**

IF SWALLOWED—Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

IF INHALED—Remove victim to fresh air and apply respiration if indicated.

IF ON SKIN—Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES—Rinse eyes for at least 15 minutes with water and call a physician immediately.

NOTE TO PHYSICIAN: 2-PAM is antidotal and may be administered in conjunction with atropine sulfate.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 119-14
EPA Est. No. 1155-011

NET WEIGHT ONE POUND

Manufactured By
SEACOAST LABORATORIES, INC.
EAST BRUNSWICK, N.J. 08816

DIRECTIONS CONTINUED

Use a hand operated duster to apply this product.

FOLIAR APPLICATION

VEGETABLES: Use at the rate of 1 1/2 pounds per 1000 square feet, except as noted. Repeat applications when necessary.

BROCCOLI, CAULIFLOWER, CABBAGE: Aphids, Diamond-back Moths and Imported Cabbageworms. Do not apply to Broccoli and Cauliflower within 5 days of harvest; Cabbage within 7 days of harvest.

CELERY: Aphids and Flea Beetles. Use 1/2 pound per 1000 square feet. Do not apply within 10 days of harvest.

CUCUMBERS: Aphids, Serpentine Leafminers and Thrips. Do not apply within 7 days of harvest.

LETTUCE: Aphids and Leafminers. Use 1/2 pound per 1000 square feet. Do not apply within 10 days of harvest.

ONIONS: Onion Thrips. Do not apply in onions within 10 days of harvest.

PEPPERS: Aphids. Use 1 1/4 pound per 1000 square feet. Do not apply within 5 days of harvest.

POTATOES: Aphids, Flea Beetles, Colorado Potato Beetles, Leaf Miners. Do not apply within 35 days of harvest.

SNAP BEANS: Black Bean Aphids, Mexican Bean Beetles. Do not apply later than 7 days before harvest.

TOMATOES: Aphids, Leaf Miners, Vinegar Flies. Do not apply within 3 days of harvest.

SOIL APPLICATION

To control CUTWORMS, GARDEN SYMPHYLAN, CRUICKETS, ROOT MAGGOTS, and WIREWORMS prior to planting of the following vegetables: Beans, Broccoli, Cabbage, Carrots, Cauliflower, Corn, Cucumbers, Endive, Lettuce, Melons, Onions, Parsnips, Peppers, Squash and Tomatoes. Apply evenly as a broadcast treatment at the rate of 1 pound per 225 square feet. Immediately after application, mix thoroughly in upper 6-8 inches of soil.

WARRANTY STATEMENT

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

ACCEPTED

JUL 23 1980

PRECAUTIONARY STATEMENTS

Hazards to Humans
(and Domestic Animals)

CAUTION

AVOID BREATHING OF DUST, AVOID CONTACT WITH SKIN, EYES AND CLOTHING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. WASH THOROUGHLY AFTER HANDLING. HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL THE FOLLOWING AFTERNOON.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE—Keep out of reach of children, pets and other animals. Avoid moisture. Avoid contamination of feed or foodstuffs by storage or disposal. Open burning and dumping prohibited. Do not reuse empty container.

DISPOSAL—Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State, or Local disposal authorities.

CONTAINER DISPOSAL—Do not reuse empty container. Wrap container and put in trash collection.

NOT FOR USE IN THE HOME



***DIAZINON®**
2% DUST

**For Control of Certain INSECTS
That Infest VEGETABLES**

ACTIVE INGREDIENTS: O, O-diethyl O- (2-isopropyl 6-methyl-4-pyrimidinyl) phosphorothioate 2.00%
INERT INGREDIENTS 98.00%

Diazinon® Trademark of CIBA-CALCI Corp.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**STATEMENT OF
PRACTICAL TREATMENT**

IF SWALLOWED—Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

IF INHALED—Remove victim to fresh air and apply respiration if indicated.

IF ON SKIN—Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES—Rinse eyes for at least 15 minutes with water and call a physician immediately.

NOTE TO PHYSICIAN: 2-PAM is antidotal and may be administered in conjunction with atropine sulfate.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 1159-124
EPA Est. No. 1159-N1

NET WEIGHT ONE POUND

Manufactured By
SEACOAST LABORATORIES, INC.

EAST BRUNSWICK, N.J. 08816

1159-124

PM 15 10F1

DIRECTIONS CONTINUED

Use a hand operated duster to apply this product.

FOLIAR APPLICATION

VEGETABLES: Use at the rate of 1 1/2 pounds per 1000 square feet, except as noted. Repeat applications when necessary.

BROCCOLI, CAULIFLOWER, CABBAGE: Aphids, Diamond-back Moths and Imported Cabbageworms. Do not apply Broccoli and Cauliflower within 5 days of harvest. Cabbage within 7 days of harvest.

CELERY: Aphids and Flea Beetles. Use 1/2 pound per 1000 square feet. Do not apply within 10 days of harvest.

CUCUMBERS: Aphids, Serpentine Leafminers and Thrips. Do not apply within 7 days of harvest.

LETTUCE: Aphids and Leafminers. Use 1/2 pound per 1000 square feet. Do not apply within 10 days of harvest.

ONIONS: Onion Thrips. Do not apply in onions within 10 days of harvest.

PEPPERS: Aphids. Use 1 1/4 pound per 1000 square feet. Do not apply within 5 days of harvest.

POTATOES: Aphids, Flea Beetles, Colorado Potato Beetle and Leaf Miners. Do not apply within 35 days of harvest.

SNAP BEANS: Black Bean Aphids, Mexican Bean Beetle. Do not apply later than 7 days before harvest.

TOMATOES: Aphids, Leaf Miners, Vinegar Flies. Do not apply within 3 days of harvest.

SOIL APPLICATION

To control CUTWORMS, GARDEN SYMPHYLAN, CRUCIFEROUS ROOT MAGGOTS, and WIREWORMS prior to planting the following vegetables: Beans, Broccoli, Cabbage, Cauliflower, Corn, Cucumbers, Endive, Lettuce, Melon, Parsnips, Peppers, Squash and Tomatoes. Apply evenly broadcast treatment at the rate of 1 pound per 225 square feet. Immediately after application, mix thoroughly in upper 6 inches of soil.

WARRANTY STATEMENT

Seller makes no warranty, expressed or implied, that the use of this product other than indicated on the label assumes all risk of use and/or handling of this material such use and/or handling is contrary to label instructions.

ACCEPTED

JUL 23 1980

PRECAUTIONARY STATEMENTS

Hazards to Humans
(and Domestic Animals)

CAUTION

AVOID BREATHING OF DUST, AVOID CONTACT WITH SKIN, EYES AND CLOTHING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. WASH THOROUGHLY AFTER HANDLING. HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL THE FOLLOWING AFTERNOON.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crops, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

Directions for Use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE—Keep out of reach of children, pets and other animals. Avoid moisture. Avoid contamination of feed or foodstuffs by storage or disposal. Open burning and dumping prohibited. Do not reuse empty containers.

DISPOSAL—Pesticide spray mixture or residue that is not to be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State, or local disposal authorities.

CONTAINER DISPOSAL—Do not reuse empty container. Wrap container and put in trash collection.

NOT FOR USE IN THE HOME



***DIAZINON®**
2% DUST

**For Control of Certain INSECTS
That Infest VEGETABLES**

ACTIVE INGREDIENTS: O, O-diethyl O-(2-isopropyl-5-methyl-4-pyrimidinyl) phosphorothioate 1.00%
INERT INGREDIENTS 0.00%
© 1979, a Division of CIBA (Gen) Corp.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**STATEMENT OF
PRACTICAL TREATMENT**

IF SWALLOWED—Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

IF INHALED—Remove victim to fresh air and apply respiration if indicated.

IF ON SKIN—Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES—Rinse eyes for at least 15 minutes with water and call a physician immediately.

NOTE TO PHYSICIAN: 2-PAA is anticholinergic and may be administered in conjunction with atropine sulfate.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

174-47g (1.00000)
174-47 (1.00000)

NET WEIGHT ONE POUND

Manufactured By
SEACOAST LABORATORIES, INC.

EAST BRUNSWICK, N.J. 08816

1159-124

PM 15 10F1

DIRECTIONS CONTINUED

Use a hand operated duster to apply this product.

FOLIAR APPLICATION

VEGETABLES: Use at the rate of 1 1/2 pounds per 1000 square feet, except as noted. Repeat applications when necessary.

BROCCOLI, CAULIFLOWER, CABBAGE: Aphids, Diamond-back Moths and Imported Cabbageworms. Do not apply to Broccoli and Cauliflower within 5 days of harvest. Cabbage within 7 days of harvest.

CELERY: Aphids and flea Beetles. Use 1/2 pound per 1000 square feet. Do not apply within 10 days of harvest.

CUCUMBERS: Aphids, Serpentine Leafminers and Thrips. Do not apply within 7 days of harvest.

LETTUCE: Aphids and Leafminers. Use 1/2 pound per 1000 square feet. Do not apply within 10 days of harvest.

ONIONS: Onion Thrips. Do not apply in onions within 10 days of harvest.

PEPPERS: Aphids. Use 1 1/2 pound per 1000 square feet. Do not apply within 5 days of harvest.

POTATOES: Aphids, Flea Beetles, Colorado Potato Beetle, Leaf Miners. Do not apply within 35 days of harvest.

SNAP BEANS: Black Bean Aphids, Mexican Bean Beetle. Do not apply later than 7 days before harvest.

TOMATOES: Aphids, Leaf Miners, Vinegar Flies. Do not apply within 3 days of harvest.

SOIL APPLICATION

To control CUTWORMS, GARDEN SYMPHYTAN, LEECHES, ROOT MAGGOTS, and WIREWORMS, prior to planting of the following vegetables: Beans, Broccoli, Cabbage, Cauliflower, Corn, Cucumbers, Endive, Lettuce, Melon, Parsnips, Peppers, Squash and Tomatoes. Apply with broadcast treatment at the rate of 1 pound per 200 square feet. Immediately after application, mix thoroughly in soil 2-3 inches or so.

WARRANTY STATEMENT

Seller makes no warranty, expressed or implied, that the use of this product other than as indicated on the label assures all risk of use and the liability of the user for any such use and/or handling, or for any other liability.

46