

DIRECTIONS


LAWN CHINCH BUGS: For control of chinch bugs in lawns, thoroughly wet down the grass and apply at the rate of 4 to 6 fluid ounces (8-12 tablespoons) in 12 gallons of water per 400 sq. ft. Application should be repeated in 7-10 days if necessary. The same dosage is effective for control of **CLOVER MITES**.

ORNAMENTALS: For control of those insects listed below attacking arbor vitae, azaleas, carnations, chrysanthemums, gladoli, holly, juniper, rhododendron, roses, pine, spruce, maple and other flowers, shrubs and shade trees mix 2½ fluid ounces (5 tablespoons) in 3 gallons of water and apply as a thorough foliar spray using a knapsack, hose end or hand type sprayer. Treatment should be made when insects first appear and repeated as infestations warrant. (NOTE: Do not use on certain ferns, poinsetta, hibiscus, papaya, pilea and gardenia, since injury may occur.)
Aphids, Bagworms, Flea Beetles, Carnation Bud mites, Thrips, Holly Bud Moth, Dipterous Leaf Miners, Leafhoppers, White Flies, Two-spotted mites, Cyclamen mites, Clover mites, Privet mites, European Red Mites, Scale crawlers: (Cottony Cushion, Lecanium, Pine Needle, San Jose and Soft Scales)

FRUITS and VEGETABLES: For control of those insects listed below use 2½ fluid ounces (5 tablespoons) in 3 gallons of water and apply as a thorough cover spray. Repeat sprays at 7-10 day intervals or as infestations warrant. Allow interval indicated between last spray application and picking of fruits or vegetables.

| Crop | Insects | Interval Between Last Spray & Picking (Days) |
|----------------|--|--|
| Apple and Pear | Codling Moth, Apple Maggot, Aphids, Pear Psylla, Mites, San Jose and Forbes Crawlers | 14 |
| Cherry | Black Cherry Aphid, Cherry Fruit Fly, San Jose Scale Crawlers | 10 |

SEACOAST LABORATORIES, INC.
EAST BRUNSWICK, N. J.



**DIAZINON
SPRAY
For LAWN
CHINCH BUGS**

**CLOVER MITES and Many INSECTS
on ORNAMENTALS —
FRUITS and VEGETABLES**

USDA Reg. No. 1159-125

WARNING

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:

| | |
|--|---------------|
| O, O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate | 10.0% |
| Aromatic petroleum derivative solvent | 82.5% |
| INERT INGREDIENTS | 07.5% |
| Total | 100.0% |

NET CONTENTS — 1 PINT

| Crop | Insects | Interval Between Last Spray & Picking (Days) |
|---------------------------|---|--|
| Grape | Mites, Leaf Hoppers, Grape Berry Moth, Grape Leaf Folder | 18 |
| Peaches | Aphids | 20 |
| Apricots | Mites | 10 |
| Nectarines | San Jose Scale Crawlers | 10 |
| Snap Beans and Lima Beans | Black Bean Aphid, Mexican Bean Beetle, Mites, Leaf Hoppers, Dipterous Leaf Miners | 7 |
| Plums | Mites | |
| Prunes | Aphids | 10 |
| Cantaloupes | Aphids | 3 |
| Watermelons | Leafhoppers | 3 |
| Squash | Mites | 7 |
| Cucumbers | Aphids | 7 |
| Cabbage | Aphids, Diamondback Moth, Imported Cabbage Worm | 7 |
| Lettuce | Aphids, Dipterous Leaf Miners | 10 |
| Peppers | Aphids | 5 |
| Potatoes | Aphids, Flea Beetles, Colorado Potato Beetle, Dipterous Leaf Miners | 14 |
| Tomatoes | Aphids, Dipterous Leaf Miners, Vinegar Flies | 3 |

WARNING: May be fatal if swallowed. Do not breathe spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of children and domestic animals. May be absorbed through skin. Do not store or use near heat or open flame. Contact with skin can cause Toxic Symptoms. Toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers.

A.C. ...
Sep 19-1974
1159-125

1001
 pm 15
 1159-125

DIRECTIONS

LAWN CHINCH BUGS: For control of chinch bugs in lawns, thoroughly wet down the grass and apply at the rate of 4 to 8 fluid ounces (8-12 tablespoons) in 12 gallons of water per 400 sq. ft. Application should be repeated in 7-10 days if necessary. The same dosage is effective for control of CLOVER MITES.

ORNAMENTALS: For control of those insects listed below attacking arbor vitae, azaleas, carnations, chrysanthemums, gladiol, holly, juniper, rhododendron, roses, pine, spruce, maple and other flowers, shrubs and shade trees mix 2 1/2 fluid ounces (5 tablespoonsful) in 3 gallons of water and apply as a thorough foliar spray using a knapsack, hose end or hand type sprayer. Treatment should be made when insects first appear and repeated as infestations warrant. (NOTE: Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia, since injury may occur.) Aphids, Bagworms, Flea Beetles, Carnation Bud mites, Thrips, Holly Bud Moth, Dipterous Leaf Miners, Leafhoppers, White Flies, Two-spotted mites, Cyclamen mites, Clover mites, Privet mites, European Bed Mites, Saw's crawlers (Cottony Cushion, Lecanium, Pine Needle, San Jose and Soft Scales).

FRUITS and VEGETABLES: For control of those insects listed below use 2 1/2 fluid ounces (5 tablespoonsful) in 3 gallons of water and apply as a thorough cover spray. Repeat sprays at 7-10 day intervals or as infestations warrant. Allow interval indicated between last spray application and picking of fruits or vegetables.

| Crop | Insects | Interval Between Last Spray & Picking (Days) |
|----------------|--|--|
| Apple and Pear | Codling Moth, Apple Maggot, Aphids, Pear Psylla, Mites, San Jose and Forbes Crawlers | 14 |
| Cherry | Black Cherry Aphid, Cherry Fruit Fly, San Jose Scale Crawlers | 10 |

SEACOAST LABORATORIES, INC.
 EAST BRUNSWICK, N. J.



**DIAZINON
 SPRAY
 For LAWN
 CHINCH BUGS**

**CLOVER MITES and Many INSECTS
 on ORNAMENTALS —
 FRUITS and VEGETABLES**

WARNING
 KEEP OUT OF REACH OF CHILDREN
 SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:
 O,O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate 10.0%
 Aromatic petroleum derivative solvent 82.5%
 INERT INGREDIENTS 07.5%
 Total 100.0%

NET CONTENTS — 1 PINT

| Crop | Insects | Interval Between Last Spray & Picking (Days) |
|---------------------------|---|--|
| Grape | Mites, Leaf Hoppers, Grape Berry Moth, Grape Leaf Folder | 18 |
| Peaches | Aphids | 20 |
| Apricots | Mites | 10 |
| Nectarines | San Jose Scale Crawlers | 10 |
| Snap Beans and Lima Beans | Black Bean Aphid, Mexican Bean Beetle, Mites, Leaf Hoppers, Dipterous Leaf Miners | 7 |
| Plums | Mites | 10 |
| Prunes | Aphids | 10 |
| Cantaloupes | Aphids | 3 |
| Watermelons | Leafhoppers | 3 |
| Squash | Mites | 7 |
| Cucumbers | Aphids | 7 |
| Cabbage | Aphids, Diamondback Moth, Imported Cabbage Worm | 7 |
| Lettuce | Aphids, Dipterous Leaf Miners | 10 |
| Peppers | Aphids | 5 |
| Potatoes | Aphids, Flea Beetles, Colorado Potato Beetle, Dipterous Leaf Miners | 14 |
| Tomatoes | Aphids, Dipterous Leaf Miners, Vinegar Flies | 8 |

WARNING: May be fatal if swallowed. Do not breathe spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of children and domestic animals. May be absorbed through skin. Do not store or use near heat or open flame. Contact with skin can cause Toxic Symptoms. Toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers.

ACCEPTED
 SEP 19-1974
 1159-125