

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Keep children and pets off treated areas until surfaces have dried.

Kitchen utensils such as teaspoons should not be used for food purposes after use in measuring pesticide.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply where runoff is likely to occur. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE. POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. MINIMUM FLASH POINT 100°F

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

LAWN CHINCH BUGS For control of chinch bugs in lawns, thoroughly wet down the grass and apply at the rate of 4 to 6 fluid ounces (8-12 tablespoons) in 2 gallons of water per 400 sq. ft. Application should be repeated in 7-10 days if necessary.

The same dosage is effective for control of CLOVER MITES.

ORNAMENTALS For control of those insects listed below attacking arbor vitae, azaleas, carnations, chrysanthemums, gladioli, holly, juniper, rhododendron, roses, pine, spruce, maple and other flowers, shrubs and shade trees mix 2 1/2 fluid ounces (5 tablespoons) in 3 gallons of water and apply as a thorough foliar spray using a knapsack, hose end or hand type sprayer. Treatment should be made when insects first appear and repeated as infestations warrant. (NOTE: Do not use on certain ferns, *Ponsetia*, *Hibiscus*, *Papaya*, *Pilea* and *Gardenia* since injury may occur.) Aphids, Bagworms, Flea Beetles, Carnation Bud mites, Thrips, Holly Bud Moth, Dipterous Leaf Miners, Leafhoppers, White Flies, Two-spotted mites, Cyclamen mites, Clover mites, Pinetree mites, European Red Mites, Scale Crawlers (Cottony Cushion, Lecanium, Pine Needle, San Jose and Soft Scale).

FRUIT and VEGETABLES For control of those insects listed below use 2 1/2 fluid ounces (5 tablespoons) in 3 gallons of water and apply as a thorough cover spray. Repeat sprays at 7-10 day intervals or as infestations warrant. Allow interval indicated between last spray application and picking of fruits or vegetables.



CHINCH BUG

FOR CLOVER MITES and MANY INSECTS ON LAWNS
ORNAMENTALS - FRUITS and VEGETABLES

ACTIVE INGREDIENTS

Diazinon 60-0-ethyl 0-0-propyl 4-methyl 10.00%
Diethylthiophosphorothioate 77.00%
Aromatic petroleum distillate 12.91%
INERT INGREDIENTS
U.S. Patent No. 2,754,743
Diazinon, Trademark of CIBA-Geigy Corp.

**KEEP OUT OF REACH OF CHILDREN.
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS**

If swallowed - Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.

If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin - Remove contaminated clothing and wash affected areas with soap and water.

If in eyes - Flush eyes with plenty of water. Call a physician immediately.

Note to Physician - Solvent prevents aspiration hazard. Gastric lavage may be indicated if material was taken internally.

EPA Reg No 1159-125 EPA Est No 1159-N21

NET CONTENTS — ONE QUART



ACCEPTED
with COMMENTS
in EPA Letter Dated:

SEP 26 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
1159-125

Crop	Insects	Interval Between Last & Picking
Apple and	Coding Moth, Apple Margot Aphids, Pear Psylla, Mites San Jose and Forbs Crawlers	
Cherry	Black Cherry Aphid Cherry Fruit Fly San Jose Scale Crawlers	
Grape	Mites, Leaf Hoppers Grape Berry Moth Grape Leaf Folder	
Peaches	Aphids	
Apricots	Mites	
Nectarines	San Jose Scale Crawlers	
Snap Beans and Lima Beans	Black Bean Aphid Mexican Bean Beetle Mites Leaf Hoppers Dipterous Leaf Miners	
Plums	Mites	
Prunes	Aphids	
Cantaloupes	Aphids	
Watermelons	Leafhoppers	
Squash	Mites	
Cucumbers	Aphids	
Cabbage	Aphids, Diamondback Moth Imported Cabbage Worm	
Lettuce	Aphids Dipterous Leaf Miners	
Peppers	Aphids	
Potatoes	Aphids, Flea Beetles Colorado Potato Beetle Dipterous Leaf Miners	
Tomatoes	Aphids Dipterous Leaf Miners Vinegar Flies	

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by stor disposal. Open dumping is prohibited.

STORAGE: Keep out of reach of children, pets and other animals. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide waste toxic.

Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by us according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap and dispose in collection.

NOTICE TO BUYER

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or harm.

1159-125