



# TWIN LIGHT

## SEVIN<sup>★</sup>

### 5% DUST 5%

**ACTIVE INGREDIENTS:**  
Carbaryl 1-naphthyl N-methylcarbamate 5.00%

**INERT INGREDIENTS:** 95.00%

## FOR JAPANESE BEETLES

**AND CERTAIN INSECTS THAT INFEST ORNAMENTALS - VEGETABLES**

\*Trade Mark of Union Carbide Corporation for 1-Naphthyl N-Methylcarbamate

**NET WEIGHT FOUR POUNDS**

USDA Reg No. 1159-128

### CAUTION

**KEEP OUT OF REACH OF CHILDREN**  
**SEE REAR PANEL FOR COMPLETE CAUTION STATEMENT**




#### DIRECTIONS

**ORNAMENTALS:** For Japanese Beetle, Bagworm, Boxwood Leaf Miner, Flea Beetle, Lace Bugs, Leafhoppers, Leaf Rollers, Periodical Cicada, Rose Aphid, Rose Slug, Scale Insects and Thrips. Apply when insects appear and repeat as needed.

**NOTE:** For Optimum Scale control apply when crawlers are present.

**VEGETABLES: BEANS** — Mexican Bean Beetle, Leafhoppers, Lygus and Stink Bugs, use ½ to 1 pound per 1000 square feet. Bean Leaf Beetle, Cucumber Beetle, Flea Beetle and Japanese Beetle, use 1 pound per 1000 square feet.

**CUCUMBER and SUMMER SQUASH:** Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs and Flea Beetles, use ½ pound per 1000 square feet.

**TOMATO, EGGPLANT and PEPPER:** Tomato Fruitworm, Fall Armyworm, Flea Beetle and Leafhoppers, use ½ to 1 pound per 1000 square feet.

Tomato Hornworm, European Corn Borer and Stink Bugs, use ½ to 1 pound per 1000 square feet.

**CORN:** Corn Earworm, European Corn Borers, Fall Armyworm, Sap Beetles and Japanese Beetles, use 1 pound per 1000 square feet.

**NOTE:** For corn earworm on corn make 3 or more applications to the silks at two to three day intervals, starting when silks first appear. For European corn borers make thorough application into the whorls when plant is knee high and repeat at weekly intervals until ears are ready for picking.

**POTATO:** Colorado Potato Beetles, Leafhoppers and Flea Beetles, use ½ to 1 pound per 1000 square feet.

**NOTE:** Do not use on Boston Ivy.

#### CAUTION

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid breathing dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

**KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

Do not feed treated bean vines to meat and dairy animals.

May be applied up to and including the day of harvest on crops listed above for human consumption.

Do not apply within 7 days of harvest for forage.

Do not reuse container. Destroy when empty.

To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material.