



Net Weight  
25 pounds

# SEVIN\*

## 5% DUST

FOR

# JAPANESE BEETLES

AND CERTAIN INSECTS THAT INFEST  
ORNAMENTALS — VEGETABLES

\* Trade Mark of Union Carbide Corporation for 1-Naphthyl N-Methylcarbamate

#### ACTIVE INGREDIENTS.

Carbaryl 1-naphthyl N-methylcarbamate . . . . . 5.00%

INERT INGREDIENTS . . . . . 95.00%

EPA Reg. No. 1159 - 128

EPA Est. No. 1159-NJ-1

KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED**—Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED**—Remove victim to fresh air and apply respiration if indicated.

**IF ON SKIN**—Wash promptly with soap and water. Rinse thoroughly.

**IF IN EYES**—Rinse eyes for at least 15 minutes with water and call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS.

ACCEPTED  
APR 1 8 1966  
U.S. DEPARTMENT OF AGRICULTURE  
Plant Industry Administration  
Washington, D.C. 20250  
EPA Reg. No. 1159-128

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels

#### DIRECTIONS

Apply only as specified on this label

**ORNAMENTALS:** For Japanese Beetle, Bagworm, Boxwood Leaf Miner, Flea Beetle, Lace Bugs, Leafhoppers, Leaf Rollers, Periodical Cicada, Rose A, Mid, Rose Slug, Scale Insects and Thrips. Apply when insects appear and repeat as needed.

**NOTE:** For Optimum Scale control apply when crawlers are present

**VEGETABLES: BEANS**—Mexican Bean Beetle, Leafhoppers, Lygus and Stink Bugs, use ½ to 1 pound per 1000 square feet. Bean Leaf Beetle, Cucumber Beetle, Flea Beetle and Japanese Beetle use 1 pound per 1000 square feet.

**CUCUMBER and SUMMER SQUASH:** Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs and Flea Beetles, use ½ pound per 1000 square feet.

**TOMATO, EGGPLANT and PEPPER:** Tomato Fruitworm, Fall Armyworm, Flea Beetle and Leafhoppers, use ½ to 1 pound per 1000 square feet.

**Tomato Hornworm, European Corn Borer and Stink Bugs,** use ½ to 1 pound per 1000 square feet.

**CORN:** Corn Earworm, European Corn Borers, Fall Armyworm, Sap Beetles and Japanese Beetles, use 1 pound per 1000 square feet.

**NOTE:** For corn earworm on corn make 3 or more applications to the silks at two to three day intervals starting when silks first appear. For European corn borers make thorough application into the whorls when plant is knee high and repeat at weekly intervals until ears are ready for picking.

**POTATO:** Colorado Potato Beetles, Leafhoppers and Flea Beetles, use ½ to 1 pound per 1000 square feet.

**NOTE:** Do not use on Boston Ivy. **CAUTION**

#### KEEP OUT OF REACH OF CHILDREN AND ANIMALS

May be harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not feed treated bean vines to meat and dairy animals.

May be applied up to and including the day of harvest on crops listed above for human consumption. Do not apply within 7 days of harvest for forage.

#### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply when conditions favor runoff from area treated. Do not apply when weather conditions favor drift from areas treated.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Keep out of reach of children, pets and other animals. Avoid moisture.

**PESTICIDE DISPOSAL:** Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

**CONTAINER DISPOSAL:** Do not reuse empty container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Securely wrap original container in several layers of newspaper and discard in trash.

**Seacoast**  
LABORATORIES, INC.