

[1159-135]



Manufactured by  
**SEACOAST LABORATORIES INC.**  
EAST BRUNSWICK, N. J. 08816

9,30,70

11/9/75

**ACTIVE INGREDIENTS:**

*Powdered Stem of Ryania Speciosa	5.00%
Pyrethrins	.10%
**Technical Piperonyl Butoxide	1.00%
Rotenone	1.00%
Other Cube Extracts	1.50%

**INERT INGREDIENTS:** 91.40%

\*Equivalent to .011 Ryanodine

\*\*Equivalent to .8% (butylcarbityl) (6-propylpiperonyl) ether and .2% related compounds.

**DIRECTIONS**

USDA Reg. No. 1159-135

Apply at the rate of 1/2 pound to 1 pound per 1000 square feet of vegetable gardens except as noted on this label.

For best results plants should be dusted or sprayed as soon as pests appear and repeated at 7 day intervals. Both lower and upper sides of plant leaves should be thoroughly dusted.

**SPRAYING DILUTIONS:** Use 6 - 7 level tablespoonfuls per gallon of water. Agitate well. Apply this dilution at the rate of 5 gallons per 1000 square feet of vegetable gardens. Shake sprayer often while spraying. On ornamentals and fruits be sure to cover upper and lower leaves thoroughly.

**FRUIT TREES:** insects Controlled - Japanese Beetles and non-wooly Aphids on Apples, Peaches, Cherries and Grapes. Fruit Flies on Cherries; Pear Psyla on Pears.

**SMALL FRUITS:** Leaf Beetle and Spittlebug on Strawberries; Maggot on Blueberries; Aphids on Blackberries; Fruitworm and Sowlly on Raspberries.

**NOTE:** Cover upper and lower leaf sides thoroughly throughout trees.

**VEGETABLES:** Young Tomato Hornworm and Aphids on Tomatoes, Peppers and Eggplants. Bean Leafhoppers and Mexican Bean Beetles on Beans; Cucumber Beetle and Aphids on Beans, Cucumbers, Melons, Squash and Peas. Flea Beetle on Potatoes, Peppers, Eggplants, Sweet Potatoes; Colorado Potato Beetle on Potatoes and Tomatoes. Cabbage Loopers, Diamond Back Moth and 6-Spotted Leafhoppers on Cabbage, Broccoli, Cauliflower and Lettuce.

**Corn Borer on Corn -** Use 8 pounds per 1000 square feet. Applications should begin when eggs begin to hatch and should be repeated at 5 to 7 day intervals.

**NOTE:** Do not make any application closer than 1 day to harvest on edible crops

**ORNAMENTALS:** Lacebugs and Aphids on Rhododendrons, Azaleas and Andromeda

**ROSES and FLOWERS:** Aphids, Thrips, Whitefly and Japanese Beetles.

**CAUTION** Avoid inhalation of dust or spray mist. Avoid contamination of feed and foodstuffs.

**FISH AND WILDLIFE CAUTION:** This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.