




GRANULAR LAWN FUNGICIDE



For Control of: **BROWN PATCH
COPPER, DOLLAR and
LEAF SPOT • SNOW MOLD**

ACTIVE INGREDIENT:

Thiram (Tetramethylthiuram Disulfide)

5.00%

*Cadmium Chloride

.75%

INERT INGREDIENTS:

94.25%

*Total Cadmium

.43%

EPA Reg.
No. 1159-136

**APPLY with SPREADER
COVERS 5,000
SQUARE FEET**

For the prevention and control of Copper Spot, Dollar Spot, Leaf Spot (Helminthosporium-Curvularia), Brown Patch and Snow Mold on Lawns containing Kentucky Blue including Merion and other strains, the various Fescues, Bents and most other grasses.

SPECIAL INSTRUCTIONS

Do not water or mow lawn area for 24 hours after application. Application will be more effective if made in the early morning when dew is on the grass. Repeat treatments are necessary until fungus diseases disappear. Apply Contents of this bag to 5,000 square feet for prevention program and at double rate (2,500 square feet) if fungus diseases have already infested lawn area.

SPRING PROGRAM:

Application should be made immediately at a double rate (contents of this bag applied to 2,500 square feet) if no disease treatments have been given and Dollar or Leaf Spot have invaded lawn area. For a prevention program against these diseases apply TWIN LIGHT GRANULAR LAWN FUNGICIDE soon after the second mowing in the Spring and repeat every 7 to 10 days for a minimum of five times. This program will prevent Fungus activity of Dollar Spot and Leaf Spot which may kill good grasses.

SUMMER APPLICATION:

Apply immediately at double rate if Brown Patch or Curvularia Disease is active in lawn area. Prevention of these diseases may be obtained by an application of TWIN LIGHT GRANULAR LAWN FUNGICIDE every 7 to 10 days during the hot humid months.

EARLY FALL APPLICATION:

Prevent infestation of Dollar Spot and Copper Spot with treatments at 10 day intervals during the period of cooler nights.

SNOW MOLD:

Apply at double rate in the late Fall, before the first snowfall and again at double rate in January.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Do not get in eyes, on skin or on clothing. Do not breathe dust since it may prove harmful. May cause skin irritation. Wash thoroughly after handling and before eating or smoking. May be fatal if swallowed. In case of contact wash immediately with soap and water. Wash all contaminated clothing with hot water before re-use. Keep children and domestic animals off treated area.

INTERNAL: If swallowed give raw white of egg. Induce vomiting with mixture of salt, soap or mustard in warm water. Call physician at once.

This product is toxic to fish and wildlife. Keep out of lakes, streams, ponds. Do not apply where runoff is likely to occur. Do not contaminate by cleaning of equipment, or disposal of wastes. Apply this product only as directed on this label.

CONTAINER DISPOSAL:

Do not reuse empty bag. Destroy it by burying with waste or burning away from smoke or fumes.

SUGGESTED SPREADER SETTINGS

Normal Rate of 10 Pounds per 5,000 Square Feet	
Scotts	7
Lawn Beauty	6 1/2
Cyclone	5
Others	1/4 of an inch opening

The age, condition of spreader, speed of operation and roughness of lawn cause variations in the rate applied. These suggested spreader settings are for your guidance only.


Where Double Rate application is wanted, go over area twice at above settings in criss cross fashion.

NET WEIGHT 10 POUNDS

Manufactured by
SEACOAST LABORATORIES, INC.
EAST BRUNSWICK, N. J. 08816



GRANULAR LAWN FUNGICIDE



For Control of: **BROWN PATCH**
COPPER, DOLLAR and
LEAF SPOT • SNOW MOLD

ACTIVE INGREDIENT:

Thiram (Tetramethylthiuram Disulfide) 5.00%
*Cadmium Chloride .75%

INERT INGREDIENTS:

*Total Cadmium .43%

APPLY with SPREADER
COVERS 5,000
SQUARE FEET

EPA Reg.
No. 1159-136

For the prevention and control of Copper Spot, Dollar Spot, Leaf Spot (Spizidium-Curvalaria), Brown Patch and Snow Mold on Lawns containing Kentucky Blue including Merion and other strains, the various Fescues, Bents and other grasses.

INSTRUCTIONS

Water or mow lawn area for 24 hours after application. Application is effective if made in the early morning when dew is on the grass. Treatments are necessary until fungus diseases disappear. Apply Contents to 5,000 square feet for prevention program and at double rate (2,500) if fungus diseases have already infested lawn area.

PROGRAM:

Application should be made immediately at a double rate (contents of this bag to 2,500 square feet) if no disease treatments have been given and Leaf Spot have invaded lawn area. For a prevention program against Leaf Spot apply TWIN LIGHT GRANULAR LAWN FUNGICIDE soon after the spring in the Spring and repeat every 7 to 10 days for a minimum of five program will prevent Fungus activity of Dollar Spot and Leaf Spot kill good grasses.

APPLICATION:

Apply immediately at double rate if Brown Patch or Curvalaria Disease is on lawn area. Prevention of these diseases may be obtained by an application of TWIN LIGHT GRANULAR LAWN FUNGICIDE every 7 to 10 days during mid months.

EARLY FALL APPLICATION:

Prevent infestation of Dollar Spot and Copper Spot with treatments at 7 to 10 day intervals during the period of cooler nights.

SNOW MOLD:

Apply at double rate in the late Fall, before the first snowfall and repeat again at double rate in January.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Do not get in eyes, on skin or on clothing. Do not breathe dust since inhalation may prove harmful. May cause skin irritation. Wash thoroughly after handling and before eating or smoking. May be fatal if swallowed. In case of contact, wash immediately with soap and water. Wash all contaminated clothing with soap and hot water before re-use. Keep children and domestic animals off treated areas.

INTERNAL: If swallowed give raw white of egg. Induce vomiting with mixture of salt, soap or mustard in warm water. Call physician at once.

This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL:

Do not reuse empty bag. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

The age, condition of spreader, speed of operation and roughness of lawn will cause variations in the rate applied. These suggested spreader settings are for your guidance only.

Where Double Rate application is wanted, go over area twice at above settings in criss cross fashion.

NET WEIGHT 10 POUNDS

Manufactured by


SEACOAST LABORATORIES, INC.

EAST BRUNSWICK, N. J. 08816



GRANULAR FUNGICIDE

For Control of: BROWN PATCH
COPPER, DOLLAR
LEAF SPOT . S



ACTIVE INGREDIENT:

Thiram (Tetramethylthiuram Disulfide)

*Cadmium Chloride

INERT INGREDIENTS:

*Total Cadmium

EPA Reg.
No. 1159-136

For the prevention and control of Copper Spot, Dollar Spot, Leaf Spot (Helminthosporium-Curvalaria), Brown Patch and Snow Mold on Lawns containing Kentucky Blue including Merion and other strains, the various Fescues, Bents and most other grasses.

SPECIAL INSTRUCTIONS

Do not water or mow lawn area for 24 hours after application. Application will be more effective if made in the early morning when dew is on the grass. Repeat treatments are necessary until fungus diseases disappear. Apply Contents of this bag to 5,000 square feet for prevention program and at double rate (2,500 square feet) if fungus diseases have already infested lawn area.

SPRING PROGRAM:

Application should be made immediately at a double rate (contents of this bag applied to 2,500 square feet) if no disease treatments have been given and Dollar or Leaf Spot have invaded lawn area. For a prevention program against these diseases apply TWIN LIGHT GRANULAR LAWN FUNGICIDE soon after the second mowing in the Spring and repeat every 7 to 10 days for a minimum of five times. This program will prevent Fungus activity of Dollar Spot and Leaf Spot which may kill good grasses.

SUMMER APPLICATION:

Apply immediately at double rate if Brown Patch or Curvularia Disease is active in lawn area. Prevention of these diseases may be obtained by an application of TWIN LIGHT GRANULAR LAWN FUNGICIDE every 7 to 10 days during the hot humid months.

SUGGESTED SPREADER SETTINGS

Normal Rate of 10 Pounds per 5,000 Square Feet	
Scotts	7
Lawn Beauty	6 1/2
Cyclone	5
Others	1/4 of an inch opening

NET WEIGHT 10 POUNDS

Manufactured by
SEACOAST LABORATORIES
EAST BRUNSWICK, N. J.