

GRANULAR CHINCH BUG USE CONTENTS

with Diazinon*

KILLS
CHINCH BUGS

OF BAG TO COVER

UP TO 5000 SQUARE FEET

CHINCH BUGS CLOVER MITES
SOD WEBWORMS (LAWN MOTHS)
JAPANESE BEETLE GRUBS

ACTIVE HIGREDIENIS

INERT INGREDIENTS:

*O.O-diethyl O (?) isopropyl 6 methyl - I pyrimidinyl) p = shorothroate

,

1.50° / **98.50**° o

FISH AND WILDLIFE CAUTION

the second of th

DIRECTIONS - APPLY WITH SPPEADER

And the second s

.

JAPANESE BEETLE GRUBS

ACTIVE INGREDIENTS

*O,O-diethyl O-(2-isopropyl-6 methyl- 4-pyrimidinyl) phosphorothioate
!NERT INGREDIENTS:

1.50° o 98.50° o

U.S. Patent No. 2.754., 13
*Diazinon Trademark of CIBA-GEIGY Corp.

DIRECTIONS - APPLY WITH SPREADER

LAWNS: For control of CHINCH BUGS apply at the rate of 15 pounds per 2500 square feet; as soon as the pest appears. Repeat in 7 to 10 days if necessary.

Granular Chinch Bug Killer applied at the above rate also efficiently controls Sod Webworms, ants, fleas, chiggers and CLOVER MITES that infest lawns.

In the New York Metropolitan area, Chinch Bugs have one generation in June and a second in August. Consult your Extension Service Specialist as to exact timing of applications.

For control of JAPANESE BEETLE GRUBS, EUROPEAN CHAFER GR BS, and SOUTHERN CHAFER GRUBS apply at the rate of 15 pounds per 1875 square feet.

For Grub control, apply anytime between late July and early October.

For control of HYPERODES WEEVIL (New York State only), apply at the rate of 15 pounds per 2250 square feet. Treat problem areas in mid April and again in mid May.

NOTE: After application sprinkle the lawn thoroughly to wash the insecucide off the grass and on to the turf. Keep children and pets off treated area until lawn has been watered and completely dried.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid Freathing dust, Harmful it swallowed. May be absorbed through skin. Avoid storage near food and feed products. Avoid contact with skin, eyes and clothing. Avoid contamination if feed and loodstaff. Wash transporting after handling.

No not reuse empty container Destroy by his hoven, with waster burning Stovicusty from more or fame.

NET WEIGHT 30 POUNDS

SEACOAST LABORATORIES, INC.

257 HIGHWAY 18

TO BRUNSWICK TO FE Jan'6.

FISH AND WILDLIFE CAUTION

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

EPA Reg.

No 1159 145

