



# TWIN LIGHT

# ROSE

## FLOWER DUST OR SPRAY



Contains  
**MALATHION**  
**FOLPET** and  
**SEVIN\*** (Carbaryl)

**Insecticide • Fungicide • Miticide**

**ACTIVE INGREDIENTS:**

**Malathion	2.00
Folpet (N Trichloromethylthio-pyridinylamide)	1.00
Carbaryl (1 Naphthyl N-methylcarbamate)	1.00

**INERT INGREDIENTS:**

100 dimethyl dithiophosphate (100 dthp)	58.00
---	-------

Registered Trade Mark of Union Carbide Corp. for Carbaryl Insecticide

### CAUTION

**KEEP OUT OF REACH OF CHILDREN**  
**SEE REAR PANEL FOR ADDITIONAL CAUTIONS**

NET WEIGHT - 1 POUND

TWIN LIGHT ROSE and FLOWER DUST or SPRAY is an Insecticide—Fungicide—Miticide combination of MALATHION—SEVIN\* (Carbaryl)—PHALTAN (Folpet) for control of certain insects and diseases of ROSES and FLOWERS as well as certain insects that infest shrubs

\*Registered Trade Mark of Union Carbide Corporation for Carbaryl Insecticide

\*\*Phaltan is registered Trade Mark for Folpet Fungicide U. S. Pat. Nos. 2,553,770—2,553,771

To be most effective applications should be made before insects become numerous and before diseases are apparent. Protect plants through the growing season by dusting or spraying at 7 to 14 day intervals and after heavy rains. Cover foliage thoroughly but do not use in excess. Pay particular attention to cover underside of leaves

**SPRAYING**—6 to 7 level tablespoonfuls per gallon of water. One pound per 8 gallons of water. Twelve pounds per 100 gallons of water. Stir thoroughly. Pour the mixture into the sprayer and agitate while spraying in order to obtain a thorough and even coverage. In a small sprayer, shake often for agitation. A hose attached sprayer may be used

**INSECTS**—Effective for the control of the following insects: Aphids, Cicada (17 year Locusts), Clover Mites, Red Spider Mites, Japanese Beetles, Rose Chafer, Rose Leafhoppers, Leaf Rollers, Rose Slugs, Flea Beetles, Thrips, Tarnished Plant Bugs, Four Lined Leaf Bugs, Whitefly, Asiatic Garden Beetles and Black Blister Beetles on Roses, Gladiolus, Dahlias, Sweet Peas, Asters, Chrysanthemums, Delphinium, Calendula, Begonia and Snapdragon

**DISEASES**—Black Spot and Powdery Mildew of Roses; Septoria Leaf Spot of Chrysanthemums; Didymellina Leaf Spot of Iris; Rust and Alternaria Leaf Spot of Carnations; Alternaria Leaf Spot, Powdery Mildew and Anthracnose on Zinnias, Marigolds, Asters, Phlox and Snapdragons

**SHRUB INSECTS**—Aphids, Japanese Beetle, Lacebugs, Spider Mites, Azalea Scale, Scuffy Scale, Euonymus Scale, Soft Brown Scale, Boxwood Leaf Miners and Bagworms on Rhododendrum, Abies, Junipers, Arborvitae, Euonymus, Hydrangeas, Pieris, Japanese Barberry and Lilac

### CAUTION

May be harmful if swallowed. May cause irritation if dust inhaled or if eye. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Store away from children and domestic animals. Avoid contamination of feed and foodstuffs. In case of contact, flush with plenty of water. For eyes, get medical attention. Dispose of the container with care.

Do not contaminate fish ponds, streams, lakes, rivers, or other bodies of water. Do not use in or near areas where livestock, poultry, or other animals are raised. Do not use on lawns, lawns, or other areas where children play. Apply the product as specified on the label.

**NOTE FOR PHYSICIAN:** This product contains Malathion, Carbaryl, and Folpet. For further information, see the label.

Manufactured By  
**SEACOAST LABORATORIES, INC**  
EAST BRUNSWICK, N. J.

USDA Reg.  
No. 1119-111

**ACCEPTED**  
Jan. 2 1970  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT