

Just right NET WEIGHT 14 OZ.

FRUIT TREE SPRAY or DUST

USDA Reg.
No. 1159-149

DIRECTIONS

SPRAYING—Add 6 level tablespoons of Just Right Fruit Tree Spray per gallon of water (1 lb. per 10 gallons of water), stir thoroughly. Pour mixture into sprayer and agitate while spraying in order to obtain a thorough and even coverage. In a small sprayer shake often for agitation.

DUSTING—This material is ready for use and is applied by a duster. Apply the material directly on the tree. Cover the fruit and foliage thoroughly with a thin coat of dust and apply when the air is still. For timing of applications, follow directions for spraying.

Just Right FRUIT TREE SPRAY or DUST is recommended to control Woolly Apple Aphids, Green Apple Aphid, Apple Maggot, Codling Moth, European Red Mites, Clover Mites, Two Spotted Mites, Williamette Mites, Eye Spotted Bud Moth, Fruit Tree Leaf Roller, Apple Red Bug, Pear Psylla, Pear Midge, Pear Plant Bug, Tarnished Plant Bug on Apples and Pears—Plum Curculio, Oriental Fruit Moth, Cottony Peach Scale, Peach Tree Borer, Japanese Beetles on Peaches—European Red Mite on Plums and Prunes—Brown Rot on Peaches, Plums, Prunes and Cherries—Leaf Spot on Cherries and diseases such as Brooks Fruit Spot, Scab and Flyspeck on Apples.

NOTE—For Peach Tree Borer make two sprays starting about July 7-15 and another about August 1-10. Spray trunks and lower branches of trees. Consult Extension authorities for exact timing in your area. Where disease control close of harvest is necessary, apply a suitable fungicide alone in a manner which will not leave residues in excess of established tolerances. On Apples, Pears, Cherries, Plums and Prunes no closer than 3 days to harvest; 7 days before harvest on Peaches.

Grape Leafhoppers, Grape Leaf Skeletonizer, Grape Berry Moth, Japanese Beetles, Red Banded Leaf Roller, Downy Mildew and Black Rot on Grapes, Spittlebug, Leaf Roller, Strawberry Weevil on Strawberries, Blueberry Maggot, Cherry Fruitworm and Japanese Beetles on Blueberries.

NOTE—Apply as soon as insects appear and repeat as needed at 7 to 10 day intervals. Do not apply closer than 1 day before harvest on Grapes, Strawberries and Blueberries.

SUGGESTED FRUIT SPRAY and DUST SCHEDULE

DELAYED DORMANT—From first green on bud to 1/2 inch leaves on Apples, Peach and Prune.

PRE-BLOOM—From first pink to first bloom on Apples, Pear, Cherry, Peach, Plum and Prune.

PETAL FALL—When last petals have fallen on all fruits listed above.

SHUCK FALL—When half the shucks have fallen from Fruit on Cherry, Peach, Plum and Prune.

COVER SPRAYS—2 to 4 applications, begin 10 days after petal fall on Cherry, Peach, Plum & Prune.

POST HARVEST—1 or 2 applications—immediately after harvest on Cherry.

Just Right FRUIT TREE SPRAY or DUST is a combination of MALATHION, SEVIN** (Carbaryl) and CAPTAN.

ACTIVE INGREDIENTS:

*Malathion	4.00%
Captan (N-trichloromethylthio Tetrahydrophthalimide)	7.50%
Carbaryl (Naphthyl N-Methylcarbamate)	3.00%

INERT INGREDIENTS:..... 85.50%

*O,O-dimethyldithiophosphate of diethyl mercaptosuccinate
**Trade Mark of Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate.

CAUTION: Do not apply within 10 days before or after oil spray treatment. Avoid contamination of feed and foodstuffs. Skin Contact with any Insecticide may be harmful. In case of contact, flush with plenty of water; for eyes, get medical attention. Avoid unnecessary contact. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Harmful if inhaled or swallowed. Do not take internally. Avoid prolonged or repeated breathing of dust or spray. Wash hands before eating.

Note for Physician. Carbaryl is a moderate, reversible, cholinesterase inhibitor.

Do not add sulphur to the spray mixture when applying this product to Red Delicious and Baldwin Apples, as this combination with Captan may cause spotting, yellowing and dropping of leaves. Injury may occur on Bosc Pears in the northeast under certain conditions.

\$1.29

Just right

FRUIT TREE SPRAY or DUST

ACCEPTED

MAR 19 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 1159-149 SUBJECT
TO ATTACHED COMMENTS.



Insecticide • Fungicide • Miticide

CAUTION

KEEP OUT OF REACH OF CHILDREN

Read instructions

\$1.29

MSDA Reg
No. 152-149

REDUCTION

SPRAY

DUST

SPRAY or DUST

NOTE

NOTE

SUGGESTED APPLICATION SCHEDULE

BEFORE BLOSSOM
PRE-BLOSSOM
PETAL FALL
SHUCK FALL
COVER SPRAY
POST HARVEST

ACTIVE INGREDIENTS

INERT INGREDIENTS

CAUTION:

NET WT. 1.0 LB.



Insecticide • Fungicide • Miticide

ACTIVE INGREDIENTS:

CAUTION

Additional cautions