



# MALATHION

## 80% SPRAY

USDA Reg. No. 1159-X

FOR **APHIDS - MITES**  
**SCALE INSECTS** on  
**ORNAMENTAL TREES & SHRUBS**

### CAUTION

KEEP OUT OF REACH OF CHILDREN  
SEE BACK PANEL FOR OTHER CAUTIONS

#### ACTIVE INGREDIENTS:

Malathion\*.....80%

Methylated Naphthalenes... 5%

INERT INGREDIENTS.....15%

100%

\*C,O-Dimethyl Dithiophosphate of

Diethyl Mercaptosuccinate

NET CONTENTS - 1 GALLON

**POULTRY:** Chickens - Ducks - Geese - Turkeys  
**DIRECT APPLICATION:** For Northern Fowl Mite and Poultry Lice, use  $\frac{1}{2}$  tablespoonfuls per 1 gallon of water per 100 - 150 birds. Repeat in 4 to 8 weeks or when necessary.

**POULTRY HOUSES:** For Northern Fowl Mite, Chicken Red Mite and Poultry Lice, use  $\frac{1}{2}$  tablespoonfuls per 1 gallon of water. Apply spray to litter, floor space and to nests, roosts and adjacent areas, force spray into cracks and crevices. This dilution may be used as a roost paint, brush on at rate of 1 pint per 150 feet of roost. Repeat as necessary.

**NOTE:** 1) Avoid contamination of Feed, Feeding Troughs and water fountains.  
2) When spraying surfaces use a high pressure sprayer.

**ORNAMENTALS:** For control of: Soft Brown Scale, Monterey Pine Scale and Black Scale Crawlers use  $\frac{1}{2}$  pints per 100 gallons of water making one or more applications as needed.

For control of: Juniper Scale, Fletcher Scale, Bagworms and Oak Kermes use 1 pint to 100 gallons of water. Make one or more applications as needed after crawlers have settled on the foliage.

For control of: Pine Needle Scale use  $\frac{1}{2}$  pints per 100 gallons of water. Make one or more applications as needed after crawlers have settled on the foliage. For the control of: Aphids, Spider Mites, Whitefly, Mealybugs, Thrips, Japanese Beetle adults, Four-lined Leaf Bug, Tarnished Plant Bug, Rose Leafhopper and Potato Leaf Hopper use 1 pint per 100 gallons of water making one or more applications as needed.

**NOTE TO PHYSICIAN:** Malathion, upon repeated prolonged careless use, may cause cholinesterase inhibition. Atropine is antidotal.

**Note:** Do not use on Boston, Maidenhair, Pteris Ferns as well as species of Crassula. Do not apply on Petunias.

**CAUTION:** Harmful if swallowed. Avoid prolonged breathing of spray mist or dust. Avoid repeated or prolonged contact with skin. Wash thoroughly. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

**SEACOAST LABORATORIES, INC.**

257 HIGHWAY 18 • EAST BRUNSWICK, N. J. 08816

**ACCEPTED**  
10-9-67  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 159-160 SUBJECT  
TO ATTACHED COMMENTS.

8 lbs  
1 lb  
1 1/2 lbs  
3/16  
100 x 8  
500 3/2 x 8  
3/16 3/16 1/8  
14