

DUSTING: Use at the rate of 2 pounds per 1200 square feet or an area of 40×30 feet on Tomatoes, Peppers and Eggplant. Use at the rate of 1 pound per 1200 square feet on other crops listed except on cucumbers, squash, melons - use at the rate of 1 pound per 1800 square feet or an area of 30×60 teet.

SPRAYING: Mix 6 to 7 level tablespoons per gallon of water and stir thoroughly. Pour mixture into sprayer and agitate while spraying in order to obtain a thorough and even coverage. Use this dilution at the rate of 1 gallon per 100 square feet on Tomatoes, Peppers and Eggplant. Use at the rate of 1 gallon per 150 square feet on other crops except 1 gallon per 200 square feet on cucumbers, melons and squash.

TOMATOES: For control of Hornworm, Fruitworm, Flea Beetle, early and late Blight, Anthracnose, Gray and Septoria Leaf Spot.

PEPPERS and EGG PLANT: Flea Beetle, Leafhoppers, Colorado Potato Beetle, Anthracnose and Corcospora on Peppers; Phomopsis, Alternaria and Anthracnose on Egg Plant.

CUCUMBERS, SQUASH and MELQNS: Striped and Spotted Cucumber Beetle, Downy Mildew, Anthracnose, Angular Leaf Spot, Scab, Squash Bug, Flea Beetles and Leafhoppers.

NOTE: Applications may be made on above vegetables up to 1 day before harvest.

BEANS: Mexican Bean Beetle, Leafhopper, Corn Earworm, Bean Leaf Beetles, Flea Beetle, Japanese Beetle, Downy Mildew, Rust and Anthracnose.

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER. BROCCOLI and KOHLRABI: Imported Cabbage Worms, Flea Beetles, Harlequin Bug, Corn Earworm, Armyworm and Cabbage Loopers.

NOTE: Do not apply to Beans, Brussels Sprouts, Cabbage, Cauliflower, Broccoli within 7 days of harvest.

Do not apply to kohlrabi after edible parts are half grown.

SWEET CORN: For European Corn Borer dust every 5 to 7 days from the time the eggs are about to hatch, when corn is knee-high on most varieties. For Corn Earworm dust silks as soon as the silks appear. Make at least three applications at 2 to 3 day intervals.

CAUTION HARMFUL IF INHALED OR SWALLOWED. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

Avoid breathing of dust or spray. Do not take internally, SKIN CONTACT may be harmful. Avoid contact.

SWEET CORN: For European Corn Borer dust every 5 to 7 days from the time the eggs are about to hatch, when corn is knee-high on most varieties. For Corn Earworm dust silks as soon as the silks appear. Make at least three applications at 2 to 3 day intervals.

CAUTION HARMFUL IF INHALED OR SWALLOWED. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.


Avoid breathing of dust or spray. Do not take internally, SKIN CONTACT may be harmful. Avoid contact, Wash hands before eating.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of feed and foodstuffs.

Do not feed forage to animals being finished for slaughter or dairy animals. Do not reuse container. Destroy when empty.

Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.





SEACOAST LABORATORIES, INC.

257 HIGHWAY 18 . EAST BRUNSWICK, N. J. 08816