



Net Weight
25 pounds

TOMATO and VEGETABLE DUST or SPRAY

ACTIVE INGREDIENTS:

*Carbaryl (1-Naphthyl N-methylcarbamate) 5.00%
Zineb (Zinc ethylene bisdithiocarbamate) 4.50%
(Metallic Zinc content . . . 92%)

INERT INGREDIENTS 90.50%

EPA Reg. No. 1159-177

EPA Est. No. 1159-N.J.-1

Twin Light TOMATO VEGETABLE DUST or SPRAY is an insecticide-fungicide combination of SEVIN* (Carbaryl) and Zineb that controls certain insects and diseases that infest tomatoes and certain vegetables.

*Trademark of Union Carbide Corp. for (1-Naphthyl) N Methylcarbamate.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED—Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED—Remove victim to fresh air and apply respiration if indicated.

IF ON SKIN—Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES—Rinse eyes for at least 15 minutes with water and call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels

DIRECTIONS

Apply only as specified on this label.

Apply to Tomatoes and Vegetables before insects become numerous and before diseases become apparent. Protect plants through the growing season by dusting or spraying at 7 to 10 day intervals and after heavy rains. Cover foliage thoroughly but do not use in excess. Pay particular attention to cover underside of leaves.

DUSTING Use at the rate of 2 pounds per 1200 square feet or an area of 40 x 30 feet on Tomatoes, Peppers and Eggplant. Use at the rate of 1 pound per 1200 square feet on other crops listed except on cucumbers, squash, melons — use at the rate of 1 pound per 1800 square feet or an area of 30 x 80 feet.

SPRAYING Mix 6 to 7 level tablespoons per gallon of water and stir thoroughly. Pour mixture into sprayer and agitate while spraying in order to obtain a thorough and even coverage. Use this dilution at the rate of 1 gallon per 100 square feet on Tomatoes, Peppers and Eggplant. Use at the rate of 1 gallon per 150 square feet on other crops except 1 gallon per 200 square feet on cucumbers, melons and squash.

TOMATOES For control of Hornworm, Fruitworm, Flea Beetle, early and late Blight, Anthracnose, Gray and Septoria Leaf Spot.

PEPPERS and EGG PLANT Flea Beetle, Leafhoppers, Colorado Potato Beetle, Anthracnose and Corcopora on Peppers; Phomopsis Alternaria and Anthracnose on Egg Plant.

CUCUMBERS, SQUASH and MELONS Striped and Spotted Cucumber Beetle, Downy Mildew, Anthracnose, Angular Leaf Spot, Scab, Squash Bug, Flea Beetles and Leafhoppers.

NOTE: Applications may be made on above vegetables up to 1 day before harvest.

BEANS Mexican Bean Beetle, Leafhopper, Corn Earworm, Bean Leaf Beetles, Flea Beetle, Japanese Beetle, Downy Mildew, Rust and Anthracnose.

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, BROCCOLI and KOHLRABI Imported Cabbage Worms, Flea Beetles, Harlequin Bug, Corn Earworm, Armyworm and Cabbage Looper.

NOTE: Do not apply to Beans, Brussels Sprouts, Cabbage, Cauliflower, Broccoli within 7 days of harvest. Do not apply to kohlrabi after edible parts are half grown.

SWEET CORN For European Corn Borer, dust every 5 to 7 days from the time the eggs are about to hatch, when corn is knee-high on most varieties. For Corn Earworm, dust silks as soon as the silks appear. Make at least three applications at 2 to 3 day intervals.

EPA Est. No. 1159-NJ-1

EPA Reg. No. 1159-177

CAUTION

**HARMFUL IF INHALED OR SWALLOWED.
KEEP OUT OF REACH OF CHILDREN**

Avoid breathing of dust or spray. Do not take internally. **SKIN CONTACT** may be harmful. Avoid contact. Wash hands before eating.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of feed and foodstuffs.

Do not feed forage to animals being flushed for slaughter or dairy animals.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply when conditions favor runoff from area treated. Do not apply when weather conditions favor drift from areas treated.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep out of reach of children, pets and other animals. Avoid moisture.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

CONTAINER DISPOSAL: Do not reuse empty container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Securely wrap original container in several layers of newspaper and discard in trash.

Seacoast
LABORATORIES, INC.
EAST BRUNSWICK, N.J. 08816