



# CHLORDANE 45% SPRAY

**For Control of TERMITES**

EPA Reg. No. 1159-178

EPA Est. No. 1159-NJ-1

**CAUTION KEEP OUT OF  
REACH OF CHILDREN  
SEE REAR PANEL  
FOR ADDITIONAL CAUTIONS**

**DO NOT USE, POUR, SPILL OR STORE  
NEAR HEAT OR OPEN FLAME.**

**ACTIVE INGREDIENTS:**

Chlordane - Technical\* 45.0%

Petroleum Distillates 50.0%

**INERT INGREDIENTS** 5.0%

Total 100.0%


\*Contains 4 lbs. Chlordane-Tech per U.S. Gal. at 70°F

\*2,7 - Octachloro-4,7-methanotetrahydromdane and  
18 related compounds

**NET CONTENTS - ONE GALLON**

**SEACOAST LABORATORIES, INC.**

257 HIGHWAY 18 • EAST BRUNSWICK, N.J. 08816



## SUBTERRANEAN TERMITES—Directions for Non-Professional Use

Use Chlordane in a 2% water emulsion by diluting 2 quarts of Twin Light CHLORDANE 45% SPRAY with 25 Gallons of water.

### BUILDINGS HAVING BASEMENTS OR CRAWL SPACES

Dig a trench along the outside of the foundations walls. In brick or hollow block or concrete foundations, dig a trench to, **but not below**, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear feet for **each foot of depth**. A trench 3 feet deep would require 12 gallons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult the State Extension Service for advice.

**CAUTION:** This concentrate, if brought into contact with the skin can cause toxic symptoms. Avoid skin contact and inhalation. In case of spillage on skin, wash immediately with soap and water. Avoid contamination of feed, food containers and foodstuffs. Do not use on household pets or humans. Harmful if swallowed. Keep out of reach of children. Do not use this product on dairy animals, in dairy barns, or on feed or forage to be fed to dairy animals or other animals being finished for slaughter.

Do not use in poultry houses.

After the Chlordane has been applied, keep children and pets away from the treated areas until the insecticide has soaked well into the soil.

Inspect foundation annually for reinfestation.

**ENVIRONMENTAL CAUTIONS:** Toxic to fish and wildlife. Birds feeding on treated area may be killed. Do not contaminate any body of water nor apply when conditions favor runoff from are treated.

Do not contaminate water by cleaning of equipment or disposal of waste.

Do not reuse container. Destroy when empty.