



Professional

USE CONTENTS OF
BAG TO COVER UP TO
25,000 SQUARE FEET

DURSBAN®

Lawn

INSECT KILLER

ACCEPTED
AUG 25 1978
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
1159-181

EPA Reg. No. 1159-181

EPA Est. 1159-N.J.-1

FOR USE ONLY BY PROFESSIONAL TURF MAINTENANCE OPERATORS.

ACTIVE INGREDIENTS:

- Chlorpyrifos [O, O-diethyl O-(3, 5, 6-trichloro-2-pyridyl) phosphorothioate] 2.32 %
- Aromatic petroleum derivative solvent 1.30 %
- INERT INGREDIENTS: 96.38 %

* Dursban - Trademark of The Dow Chemical Co.,
U.S. Patent No. 3,244,586

(Wash granules into soil with 30 gallons of water per 1,000 sq. ft.)

**APPROXIMATE SPREADER SETTINGS
CYCLONE**

| | | |
|-------------------------|-------|---|
| Pounds per 1000 sq. ft. | | |
| 2 1/4 | 3 1/2 | 4 |

The age, condition of spreader, speed of operation and roughness of lawn will cause variations in rates applied. These suggested spreader settings are for your guidance only.

DIRECTIONS ORNAMENTAL TURF

CHINCH BUGS; SOD WEBWORMS, ANTS, EARWIGS, CRICKETS, CUTWORMS, BROWN DOG TICKS and SOWBUGS (PILLBUGS). Spread uniformly at a rate of 1 pound per 1,000 sq. ft. In northeastern states, make first application in May or June or when insects first appear. Re-treat as needed, usually in July or August

Consult your Extension Service Specialist as to exact timing of applications.

HYPERODES WEEVIL: Apply uniformly at a rate of 2 lbs. per 1,000 sq. ft. Make the first application of suspected area in mid-April followed by second application 30 days later.

JAPANESE BEETLE GRUBS and EUROPEAN CHAFER GRUBS.

Depending on infestation, apply uniformly at a rate of 2 to 4 lbs. per 1000 sq. ft. For Grub control, apply anytime between late July and early October when grubs are young and feeding near surface.

After application sprinkle the lawn thoroughly to wash the insecticide off the grass and on to the turf.

(Wash granules into soil with 30 gallons of water per 1,000 sq. ft.)

WARNING

KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed or inhaled. Do not take internally. Do not breathe dust. Wash thoroughly after handling. Do not get in eyes.

Flush contaminated eyes with plenty of water and get medical attention. Wash contaminated clothing before re-use. Keep away from foods, feedstuffs, and domestic water supplies. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

FISH AND WILDLIFE CAUTION: This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Dispose of empty container and wastes by burning or burying in non-crop lands away from domestic water supplies.

NOTE: Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

NET WEIGHT 25 POUNDS

Manufactured by

SEACOAST LABORATORIES INC.

EAST BRUNSWICK, N. J. 08816