



GRANULAR

BENOMYL TURF FUNGICIDE

Controls: DOLLAR SPOT

EPA Reg. No. 1159-183

EPA F.I. 1159-NJ-1

BROWN PATCH • STRIPE SMUT

ACTIVE INGREDIENT

Benomyl [Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate] 1.50%

INERT INGREDIENTS

98.50%

CAUTION KEEP OUT OF REACH OF CHILDREN

Avoid breathing dust. Avoid contact with skin, eyes, and clothing. May irritate nose, throat, and skin. Wash thoroughly after using. In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

Do not graze or feed clippings from treated areas to livestock. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

FISH AND WILDLIFE CAUTION

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

USE CONTENTS

**OF BAG TO COVER UP TO
3500 SQUARE FEET**

APPROXIMATE SPREADER SETTINGS

3500 Sq. Ft. 1150 Sq. Ft.

1159-183 CYCLONE	5	7
	7	9

Above spreader settings are approximate. Allowance must be made for the condition of spreader, the speed at which applied and roughness of area being treated.

DIRECTIONS — ORNAMENTAL TURF

Twin Light GRANULAR BENOMYL TURF FUNGICIDE is recommended for the control of DOLLAR SPOT, LARGE BROWN PATCH and STRIPE SMUT.

When conditions are unusually favorable for development of disease, reduce interval to 5-7 days.

BROWN PATCH • STRIPE SMUT

ACTIVE INGREDIENT

Benomyl [Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate] 1.50%

INERT INGREDIENTS

98.50%

CAUTION KEEP OUT OF REACH OF CHILDREN

Avoid breathing dust. Avoid contact with skin, eyes, and clothing. May irritate nose, throat, and skin. Wash thoroughly after using. In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

Do not graze or feed clippings from treated areas to livestock. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

FISH AND WILDLIFE CAUTION

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

USE CONTENTS

OF BAG TO COVER UP TO
3500 SQUARE FEET

APPROXIMATE SPREADER SETTINGS

3500 Sq. Ft. 1150 Sq. Ft.

CYCLONE	5	7
SCOTT	7	9

Above spreader settings are approximate. Allowance must be made for the condition of spreader, the speed at which applied and roughness of area being treated.

DIRECTIONS — ORNAMENTAL TURF

Twin Light GRANULAR BENOMYL TURF FUNGICIDE is recommended for the control of DOLLAR SPOT, LARGE BROWN PATCH and STRIPE SMUT. This product provides both an eradicant and protection action against these diseases.

For Dollar Spot (*Sclerotinia homeocarpa*), Large Brown Patch (*Rhizoctonia solani*) - Apply contents of bag, 15 pounds to 3500 square feet (35' x 100') at earliest appearance of these diseases and continue at 10 to 14 day intervals as long as needed.

When conditions are unusually favorable for development of diseases, reduce interval to 5 to 7 days.

For Stripe Smut (*Ustilago striiformis*) - Make a single application at rate of 15 pounds per 1150 square feet in October or in early spring before grass begins growth.

Note: After application, water turf sufficiently to carry product to base of plants.

NET WEIGHT 15 POUNDS

Manufactured By

SEACOAST LABORATORIES, Inc.

257 HIGHWAY 18 EAST BRUNSWICK, N.J. 08816

