

301 156107  
AUG 26 1985 17 3

Seacoast Laboratories, Inc.  
P.O. Box 157  
East Brunswick, NJ 08816

Attention: Mr. Irvin Baker

Gentlemen:

Subject: Twin Light Granular Benomyl Turf Fungicide  
EPA Registration No. 1159-183  
Your Letter Dated June 25, 1985

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted at the next printing.

1. Change "Keep out lakes, streams, or ponds" to "Do not apply directly to water."

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Henry M. Jacoby  
Product Manager (21)  
Fungicide-Herbicide Branch  
Registration Division (TS-767)

Enclosure

RD/DIS:JOB:88202:Noles:RD-50:Kendrick:898-1270:8/19/85:del.8/29/85:dej

CONCURRENCES

SYMBOL ▶								
SURNAME ▶								
DATE ▶								

**Twin<sup>®</sup>  
Light**

NET WEIGHT  
30 POUNDS



granular

# Benomyl Turf Fungicide

Controls: dollar spot  
brown patch stripe smut

Use contents of bag to cover up to,  
7,000 square feet.

ACCEPTED  
with COMMENTS  
in EPA

AUG 29 1985

Under the Fungicide, Insecticide, and Plant Growth Regulator Act of 1972, this product is registered under EPA Reg. No.

1159-183

ACTIVE INGREDIENT

EPA Reg. No. 1159-183  
EPA Est. 1159-NJ-1

Benomyl (Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate) 1.50%

INERT INGREDIENTS 98.50%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Call a physician. Do not induce vomiting unless instructed. Drink one or two glasses of water and induce vomiting by forcing back of throat with finger. Repeat until vomit is clear. Do not induce vomiting or give anything by mouth to unconscious persons.

**IF INHALED:** Remove victim to fresh air and keep respiratory tract moist.

**IF ON SKIN:** Wash promptly with soap and water. Rinse thoroughly.

**IF IN EYES:** Flush eyes for at least 15 minutes with water, wash all contact lenses.

**SEE BACK PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS**

## CONTROLS: DOLLAR SPOT • BROWN PATCH • STRIPE SMUT

*Back to back  
Reduced 35%*

### PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

#### CAUTION

Avoid breathing dust. Avoid contact with skin, eyes, and clothing. May irritate nose, throat, and skin. Wash thoroughly after using. In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention. Do not graze or feed clippings from treated areas to livestock.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.  
**STORAGE** Keep out of reach of children, pets and other animals. Avoid moisture.  
**PESTICIDE DISPOSAL** Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.  
**CONTAINER DISPOSAL** Do not reuse empty container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Wrap empty container and place in trash collection.

#### APPROXIMATE SPREADER SETTINGS

	7000 Sq. Ft.	2300 Sq. Ft.
CYCLONE . . . . .	4½	7
SCOTT (DROP) . . . .	5	9
SCOTT (BROADCAST) . . .	D½	0

Above spreader settings are approximate. Allowance must be made for the condition of spreader, the speed at which applied and the roughness of area being treated.

#### DIRECTIONS — ORNAMENTAL TURF

Twin Light GRANULAR BENOMYL TURF FUNGICIDE is recommended for the control of DOLLAR SPOT, LARGE BROWN PATCH and STRIPE SMUT. This product provides both an eradicant and protection action against these diseases.

For Dollar Spot (*Sclerotinia homoeocarpa*), Large Brown Patch (*Rhizoctonia solani*) — Apply at the rate of 30 pounds to 7000 square feet (70' x 100') at earliest appearance of these diseases and continue at 10 to 14 day intervals as long as needed.

When conditions are unusually favorable for development of disease, reduce intervals to 5 to 7 days.

For Stripe Smut (*Ustilago striiformis*) — Make a single application at rate of 30 pounds per 2300 square feet in October or in early spring before grass begins growth.

**NOTE** After application, water turf sufficiently to carry product to base of plants.

#### WARRANTY STATEMENT

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instruction.

# NET WEIGHT 30 POUNDS