



BENOMYL FUNGICIDE

**Wettable Powder For
TURF • ROSES •
FLOWERS • ORNAMENTALS**

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE REAR PANEL FOR ADDITIONAL CAUTIONS
NET WEIGHT 4 OZ.

Manufactured For
SEACOAST LABORATORIES, INC.
EAST BRUNSWICK, N. J.

ACTIVE INGREDIENT

Benomyl [Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate] 50%

INERT INGREDIENTS 50%

U. S. Pat. Pending

EPA Reg. No. 1159-184

CAUTION KEEP OUT OF REACH OF CHILDREN

MAY IRRITATE EYES, NOSE, THROAT, and SKIN

Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using. Keep away from fire or sparks. In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

IMPORTANT - Never allow Twin Light BENOMYL FUNGICIDE to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of Twin Light BENOMYL FUNGICIDE as a fungicide. Keep container closed when not in use. Do not re-use container; bury when empty.

FISH and WILDLIFE CAUTION

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS

Twin Light BENOMYL FUNGICIDE is recommended for the control of certain diseases of turf, roses flowers and ornamentals. It is highly effective for the control of many fungus diseases, providing both curative (eradicator) action and protective action. Twin Light BENOMYL FUNGICIDE is a wettable powder to be mixed with water for application as a spray. Addition of a surfactant to the spray mixture enhances curative action of the fungicide, and improves distribution of the spray on hard-to-wet plants such as roses.

TURF - Dollar Spot (Sclerotinia homecarpa), Large Brown Patch (Rhizoctonia solani): Apply at the rate of 2 oz. (12 tablespoons) per 1000 sq. ft. in sufficient water to obtain thorough coverage, usually 2 to 5 gals. per 1000 sq. ft. Apply at earliest appearance of disease and continue at 10 to 14 day

intervals as long as needed. When conditions are unusually favorable for development of disease, reduce interval to 5 to 7 days.

FUSARIUM PATCH (Fusarium nivale) - a cool season disease - Apply at the rate of 2 oz. (12 tablespoons) in 2 to 5 gals. of water per 1000 sq. ft. at the earliest appearance of disease and continue at 10 to 14 day intervals as needed.

FUSARIUM BLIGHT (Fusarium roseum and F. tricinctum) - a warm season disease - Apply at rate of 5 to 8 oz. in 2 to 5 gals. of water per 1000 sq. ft. at earliest appearance of disease and repeat 10 to 14 days later. Immediately after application, water turf sufficiently to thoroughly wet soil.

STRIPESMUT (Ustilago striiformis) - Make a single application of 6 oz. in 2 to 5 gals. of water per 1000 sq. ft. in October or in early spring before grass begins growth. After application, water turf sufficiently to carry product to base of plants.

NOTE Do not graze or feed clippings from treated areas to livestock.

ROSES (Field and Greenhouse) POWDERY MILDEW, BLACK SPOT: Use 1 tablespoonful per 2 gals. of water. Begin application when disease first appears and repeat at 10 to 14-day intervals throughout the growing season; shorten intervals during humid, rainy weather.

FLOWERS, ORNAMENTALS (Field and Greenhouse) - POWDERY MILDEW, BOTRYTIS GREY MOLD: Use 1 tablespoonful per 2 gals. of water. Begin application when disease first appears and repeat at 10 to 14-day intervals as long as needed.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and/or handling of this material when such use and/or handling is contrary to label instructions.