



*Diazinon®

TREATS 5000 SQ. FT. FOR
CHINCH BUGS. 4000 SQ. FT.
FOR JAPANESE BEETLE GRUBS

Lawn INSECT KILLER

**KILLS CHINCH BUGS CLOVER MITES
SOD WEBWORMS (LAWN MOTHS)
JAPANESE BEETLE GRUBS**

ACTIVE INGREDIENT

*O,O-diethyl O-(2-isopropyl-6 methyl-4-pyrimidinyl) phosphorothioate _____ 5%

INERT INGREDIENTS: _____ 95%

U. S. Patent No. 2,754,243

*Diazinon® Trademark of CIBA-GEIGY Corp.

DIRECTIONS — APPLY WITH SPREADER

LAWNS: For control of CHINCH BUGS apply at the rate of 2 pounds per 1000 square feet; as soon as the pest appears. Repeat in 7 to 10 days if necessary.

Diazinon 5% Granular applied at the above rate also efficiently controls Sod Webworms, ants, fleas, chiggers and CLOVER MITES that infest lawns.

In the New York Metropolitan area, Chinch Bugs have one generation in June and a second in August. Consult your Extension Service Specialist as to exact timing of applications.

For control of JAPANESE BEETLE GRUBS, EUROPEAN CHAFER GRUBS, and SOUTHERN CHAFER GRUBS apply at the rate of 2½ pounds per 1000 square feet.

For Grub control apply anytime between late July and early October. For control of HYPERODES WEEVIL (New York State only), apply at the rate of 2 pounds per 1000 square feet. Treat problem areas in mid April and again in mid May.

NOTE: After application sprinkle the lawn thoroughly to wash the insecticide off the grass and on to the turf. Keep children and pets off treated area until lawn has been watered and completely dried.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid breathing dust. Harmful if swallowed. May be absorbed through skin. Avoid storage near food and feed products. Avoid contact with skin, eyes and clothing. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling.

Do not reuse empty container. Wrap container and put in trash collection.

EPA Reg. No. 1159-188

EPA Est. No. 1159-NJ-1

APPROXIMATE SPREADER SETTINGS

Per 1000 Sq. Ft.

2 lbs. 2½ lbs.

SCOTT ___ 3½ 4

CYCLONE ___ 3 3½

The age, condition of spreader, speed of operation and roughness of lawn will cause variations in rates applied. These suggested settings are for your guidance only.

FISH AND WILDLIFE CAUTION

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Apply this product only as specified on this bag.

NET WEIGHT 10 POUNDS

SEACOAST LABORATORIES INC.

EAST BRUNSWICK, N. J. 08816