



Net Weight  
25 pounds

# MULTI-PURPOSE SPRAY

CONTROLS CERTAIN INSECTS AND DISEASES ON

**ROSES • SHRUBS  
FRUIT TREES  
VEGETABLES**

**ACTIVE INGREDIENTS**

Carbaryl (1-Naphthyl N-Methylcarbamate\*)  
Malathion\*\*  
Folpet N-(Trichloromethylthio) Phthalimide  
Kethane 1,1 Bis (Chlorophenyl) 2,2,2-Trichloroethanol\*\*\*

**INERT INGREDIENTS**

TOTAL 100.00%

\*SEVIN Reg. T.M. Carbide Corp.  
\*\*D.O. Dimethyl Dithiophosphate of diethyl mercaptosuccinate  
\*\*\*KELTHANE Reg. T.M. Rohm & Haas Co.

EPA Reg. No. 1159-197

EPA Est. No. 1159-NJ-1

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
STATEMENT OF PRACTICAL TREATMENT

Reg # 1159-197  
PM 12 p. 1-11

Subsequent planting of food and tree crops in treated soil must be made with crops on this or other carbaryl labels

**DIRECTIONS**

**DILUTION RATE** The general rate of application is 1 lb. in 20 gallons of water (3 level tablespoons per gallon) Note heavier rates required for bagworm and scale insect control

**WHEN TO APPLY** For most effective results follow a regular spray program in early Spring when insects first appear, but before diseases appear. Continue at 10 day intervals, after heavy rains, or as needed. Cover both upper and lower surfaces of foliage. For timing of application on fruit trees follow suggested schedule or consult your State Agricultural Extension Service for additional information consistent with label directions and cautions

Apply this product only as specified on this label

**RECOMMENDED USES ON ROSES, ORNAMENTALS, SHRUBS AND TREES**

**DISEASES** For the control of Black Spot and Powdery Mildew of Roses, Septoria Leaf Spot on Chrysanthemums, Rust of Carnations, Powdery Mildew and Anthracnose of Zinnias, Marigolds, Phlox, Snapdragons and Asters

**INSECTS** For the control of Aphids, Spider Mites, Japanese Beetles, Thrips, Lacebugs, Leaf Rollers, Leafminers on Boxwood, Holly and Birch, Tent Caterpillars

When used for the control of Bagworms on Spruce, Euonymus Scale Crawlers, Soft Brown Scale Crawlers, Azalea Scale Crawlers on Azaleas, Rhododendron, and other shrubs use 1 lb. in 10 gallons of water (8 level tablespoons per gallon)

**NOTE** Do not use on Boston, Maidenhair, Pteris Ferns as well as species of Crassula. Do not use on Petunias or Boston Ivy. Do not use on Japanese Maples

**RECOMMENDED USES ON FRUIT AND VEGETABLES**

**DISEASES** For the Control of Apple Scab, Sooty Blotch, Bitter Rot, Leaf Spot on Apples, Crabapples and Sour Cherries, Downy and Powdery Mildew on Cucumbers and Melons, Fruit Rot on Blackberries and Raspberries

**INSECTS** For the control of Codling Moth, Curculio, Spider Mites, Pear Psylla and Apple Maggot on APPLES. Tent Caterpillars, Black Cherry Aphid, Japanese Beetles and Two Spotted Mites on CHERRIES

Japanese Beetles, Leaf Rollers, Leafhoppers and Raspberry Aphids on BLACKBERRIES and RASPBERRIES. Japanese Beetles, Grape Berry Moth, Grape Leafhopper and Red Banded Leaf Roller on GRAPES

Aphids, Spider Mites, Flea Beetles and Colorado Potato Beetles on TOMATOES. Aphids, Cucumber Beetles and Squash Bugs on CUCUMBERS, MELONS and SQUASH

**CAUTION:**

DO NOT APPLY TO VEGETABLES WITHIN 2 DAYS OF HARVEST AND 8 DAYS ON TOMATOES.

**SUGGESTED FRUIT SPRAY SCHEDULE**

Delayed Dormant - (From first green on bud to 1/2 inch leaves on Apple)

Pre-Bloom - (From first pink to first bloom on Apple and Cherry)

Petal Fall - (When last petals have fallen) On all fruits listed above

Cover Sprays - (2 to 4 applications) Begin 10 days after petal fall on Apples

Post Harvest Spray for Cherry Leaf Spot - Apply immediately after harvest and 10 days later

**CAUTIONS: DO NOT APPLY TO FRUIT TREES WITHIN 7 DAYS OF HARVEST AND ON SOUR CHERRIES MAKE 1 APPLICATION AND THEN NOT AGAIN FOR 30 DAYS. MAY BE INJURIOUS TO SOME VARIETIES OF APPLES**

Do not apply to blackberries or raspberries within 7 days of harvest. Do not apply to grapes within 3 days of harvest

**CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust or spray mist. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

Do not store near or contaminate feed or footstuffs

Do not use on certain ferns including Boston, Maidenhair and Pteris, some species of Crassula and Boston Ivy

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply when conditions favor runoff from area treated. Do not apply when weather con-.....

ACC  
APR 18 1986  
Act  
us  
1159-197

CONTROLS CERTAIN INSECTS AND DISEASES

# ROSES • SHRUBS FRUIT TREES VEGETABLES

**ACTIVE INGREDIENTS**

Carbaryl (1-Naphthyl N-Methylcarbamate\*)

Malathion\*\*

Folpet N-(Trichloromethylthio) Phthalimide

Kelthane 1,1-Bis (Chlorophenyl) 2,2,2-Trichloroethane\*\*\*

**INERT INGREDIENTS**

ACC  
APR 18 1986  
1159-197

10.0%  
8.0%  
14.5%  
1.8%  
66.0%  
Total 100.00%

\*SEVIN Reg. T.M. Carbide Corp

\*\*D.O. Dimethyl Dithiophosphate of diethyl mercaptosuccinate

\*\*\*KELTHANE Reg. T.M. Rohm & Haas Co

EPA Reg. No. 1159-197

EPA Est. No. 1159-NL-1

## KEEP OUT OF REACH OF CHILDREN CAUTION STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED**—Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED**—Remove victim to fresh air and apply respiration if indicated.

**IF ON SKIN**—Wash promptly with soap and water. Rinse thoroughly.

**IF IN EYES**—Rinse eyes for at least 15 minutes with water and call a physician immediately.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

# Seacoast

LABORATORIES, INC.  
EAST BRUNSWICK, N.J. 08816

### SUGGESTED FRUIT SPRAY SCHEDULE

Delayed Dormant - (From first green on bud to 1/2 inch leaves on Apple)

Pre-Bloom - (From first pink to first bloom on Apple and Cherry)

Petal Fall - (When last petals have fallen) On all fruits listed above

Cover Sprays - (2 to 4 applications) Begin 10 days after petal fall on Apples

Post Harvest Spray for Cherry Leaf Spot - Apply immediately after harvest and 10 days later

**CAUTIONS** DO NOT APPLY TO FRUIT TREES WITHIN 7 DAYS OF HARVEST AND ON SOUR CHERRIES MAKE 1 APPLICATION AND THEN NOT AGAIN FOR 30 DAYS MAY BE INJURIOUS TO SOME VARIETIES OF APPLES

Do not apply to blackberries or raspberries within 7 days of harvest Do not apply to grapes within 3 days of harvest

### CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust or spray mist. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

Do not store near or contaminate feed or footstuffs

Do not use on certain ferns including Boston, Maidenhair and Pteris, some species of Crassula and Bon-in-try

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply when conditions favor runoff from area treated. Do not apply when weather conditions favor drift from areas treated.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Keep out of reach of children, pets and other animals. Avoid moisture.

**PESTICIDE DISPOSAL:** Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

**CONTAINER DISPOSAL:** Do not reuse empty container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Securely wrap original container in several layers of newspaper and discard in trash