



EPA Est. No. 1159-NJ 1

EPA Reg. No. 1159-199

**GRANULAR**

**LAWN INSECTICIDE**

**WITH**

**DURSBAN**

**USE CONTENTS OF  
BAG TO COVER UP TO  
10,000 SQUARE FEET**


**APPROXIMATE  
SPREADER SETTINGS**

Per 1000 Sq. Ft.

2 lbs. 4 lbs.

CYCLONE ..... 3 4 1/2  
SCOTT ..... 4 5 1/2

The age, condition of spreader, speed of operation and roughness of lawn will cause variations in rates applied. These suggested spread settings are for your guidance only.



**ACTIVE INGREDIENTS:**

Chlorpyrifos [O, O-diethyl O-(3, 5, 6-trichloro-2-pyridyl) phosphorothioate] ..... 1.16%  
Aromatic petroleum derivative solvent ..... .65%

**INERT INGREDIENTS:** ..... 98.19%

\* Dursban® Trademark of The Dow Chemical Co.,  
U.S. Patent No. 3,244,586

**NET WEIGHT 20 POUNDS**

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

**STATEMENT OF PRACTICAL TREATMENT**

In case of contact: for eyes, flush with water for 15 minutes; for skin, wash with plenty of soap and water. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** Active ingredient is a cholinesterase inhibitor.

Treat symptomatically. Atropine is an antidote.

**HAZARD TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

**ENVIRONMENTAL HAZARD:**

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated.

Do not allow children or pets on treated lawn until insecticide is washed off the grass into the soil and grass is completely dried.

**DIRECTIONS FOR USE  
GENERAL CLASSIFICATION**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE and DISPOSAL:**

Keep container closed.

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not reuse empty container.

Wrap container and put in trash collection.

**NOTE:** Follow directions carefully, as use, storage and handling of this material is beyond the control of the seller, the buyer assumes all risks if directions given herewith are not strictly adhered to.

**DIRECTIONS ORNAMENTAL TURF**

(Do not use on Dichondria):

**CHINCH BUGS; SOD WEBWORMS, ANTS, EARWIGS, CRICKETS, CUT-  
WORMS, BROWN DOG TICKS and SOWBUGS (PILLBUGS).**

Spread uniformly at a rate of .2 pounds per 1,000 sq. ft. In northeastern states, make first application in May or June or when insects first appear. Re-treat as needed, usually in July or August.

Consult your Extension Service Specialist as to exact timing of applications.

(Wash granules into soil with 30 gallons of water per 1,000 sq. ft.)

**HYPERODES WEEVIL:**

Apply uniformly at a rate of 4 pounds per 1,000 sq. ft. Make the first application of suspected area in mid-April followed by second application 30 days later.

To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle.

After application sprinkle the lawn thoroughly to wash the insecticide off the grass and on to the turf.

(Wash granules into soil with 30 gallons of water per 1,000 sq. ft.)

**JAPANESE BEETLE GRUBS and EUROPEAN CHAFER GRUBS:**

Apply uniformly at a rate of 4 to 8 pounds per 1,000 sq. ft.

For Grub control, apply anytime between July and early October when grubs are young and feeding near soil surface.

**NOTE:** For Hyperodes weevil and grubs make application using the above spreader settings in criss cross fashion.

After application sprinkle the lawn thoroughly to wash the insecticide off the grass and on to the turf.

(Wash granules into soil with 30 gallons of water per 1,000 sq. ft.)

Consult your Extension Service Specialist as to exact timing of applications.

**SEACOAST LABORATORIES, INC.**

**EAST BRUNSWICK, N. J. 08816**