




NET WEIGHT 40 POUNDS

BALAN 2.5% GRANULAR



ACTIVE INGREDIENT

N-butyl-N-ethyl-o,o-difluoro-2
chloro-1-methyl-4-pyridine

INERT INGREDIENTS

BALAN is a registered trademark of Seacoast Laboratories, Inc.
BENEFIN is a registered trademark of E. I. du Pont de Nemours & Co.

COVERS 1/2 ACRE

For Pre-Emergence Control of CRABGRASS and ANNUAL BLUE GRASS IN ESTABLISHED LAWNS

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED-Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
IF INHALED-Remove victim to fresh air and apply respiration if indicated.
IF ON SKIN-Wash promptly with soap and water. Rinse thoroughly.
IF IN EYES-Rinse eyes for at least 15 minutes with water and call a physician immediately.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY

STATEMENTS

EPA Est. #1159-NJ-1

EPA Reg. #1159-200

Seacoast
LABORATORIES, INC.
EAST BRIDGEWATER, N.J. 08817

Front Panel



NET WEIGHT 40 POUNDS

BALAN[®] 2.5 % GRANULAR

COVERS 1/2 ACRE

For Pre-Emergence Control of CRABGRASS and ANNUAL BLUE GRASS IN ESTABLISHED LAWNS

PRECAUTIONARY STATEMENTS

Hazards to Humans
(and Domestic Animals)

CAUTION

HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN, EYES AND CLOTHING. DO NOT APPLY BY HAND. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. WASH THOROUGHLY AFTER HANDLING AND BEFORE EATING OR SMOKING. KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL LAWN HAS BEEN WATERED AND COMPLETELY DRIED.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep out of reach of children, pets and other animals. Avoid moisture.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

CONTAINER DISPOSAL: Do not reuse empty container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Wrap empty container and place in trash collection.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

LAWN and ORNAMENTAL TURF (established turf only):

Apply at rate of 40 lbs. per 1/2 acre (21,260 sq. ft.) of lawn with a spreader for the control of Smooth and Hairy Crabgrass, Goosegrass (Silver Crabgrass), Yellow and Green Foxtail and Annual Bluegrass (Poa Annua). This product controls weeds by killing their seeds as they germinate. It does not control established weeds. Time application in the Spring before the germination of annual grass weeds tested. Water after application to wash chemical into the turf.

PRECAUTION: Do not apply in the Spring to completely new lawns started the previous Fall. Do not apply to Bentgrass putting greens or diandra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass or Centipede grass until these turfs are well established since root inhibition may result. Do not reseed within 42 days after treatment.

APPROXIMATE SPREADER SETTINGS

CYCLONE	3 1/2
SCOTT	4 1/2

The age, condition of spreader, speed of operation and roughness of lawn will cause variations in rates applied. These suggested spreader settings are for your guidance only.

Seacoast
LABORATORIES, INC.
EAST BRUNSWICK, N.J. 08816

*Asch
Paul*