



NET WEIGHT  
15 LBS. 10 OZS

1159-202  
**GARDEN WEED PREVENTER**

CONTROLS GRASSY WEEDS and most BROADLEAF WEEDS  
in SHRUB BEDS, FLOWER BEDS and VEGETABLE GARDENS

**COVERS** up to 20,000 SQ.FT. when mixed with  
the SOIL and 2,500 SQ. FT. when APPLIED  
to the SOIL SURFACE

ACTIVE INGREDIENT  
trifluralin (a,a,a-trifluoro-2,6-dinitro-N,N-  
dipropyl-p-toluidine).....1.47% ✓  
INERT INGREDIENT.....98.53%

EPA Reg. # 1159-

EPA Est. # 1159-N.J.-1

E

KEEP OUT OF REACH OF CHILDREN  
CAUTION

STATEMENT OF  
PRACTICAL TREATMENT

IF SWALLOWED - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.


✓ IF INHALED - Remove victim to fresh air and apply respiration if indicated.

IF ON SKIN - Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES - Rinse eyes for at least 15 minutes with water and call a physician immediately.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY  
STATEMENTS

IF DOCUMENT AVAILABLE  
**Seacoast**  
LABORATORIES, INC.  
EAST BRUNSWICK, N.J. 08816



E



NET WEIGHT

15 LBS. 10 OZS

## GARDEN WEED PREVENTER

### PRECAUTIONARY STATEMENTS

Hazards to Humans  
(and Domestic Animals)

#### CAUTION

HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN, EYES AND CLOTHING. DO NOT APPLY BY HAND. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. WASH THOROUGHLY AFTER HANDLING AND BEFORE EATING OR SMOKING.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Keep out of reach of children, pets and other animals. Avoid moisture.  
**PESTICIDE DISPOSAL:** Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

**CONTAINER DISPOSAL:** Do not reuse empty container. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Wrap empty container and place in trash collection.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### ESTABLISHED TREES, ORNAMENTALS, SHRUBS and FLOWERS

**APPLICATION TO SOIL SURFACE:** Where mixing Garden Weed Preventer into the soil would be impractical, application can be made to the soil surface around established trees, shrubs, flowers and roses, listed below. First remove existing weeds, then apply Garden Weed Preventer to the soil surface at the rate of 1 lb. per 160 sq. ft. Next thoroughly soak the soil to activate the herbicide.

**ESTABLISHED FLOWERS:** Recommended for use on; ACHILLEA, AGERATUM ARCTOTIS, ASTER BALSAM, CALENDULA, CALIFORNIA POPPY, CALLIOPSIS, CARNATION, CENTAUREA, CHRYSANTHEMUM, COSMOS, DAHLIA, DIMORPHETHECA, FORGET-ME-NOT, 4 O'CLOCK, GAILLARDIA, GLADIUS\*\*, HOLLYHOCK, IXORA, LOBELIA, LUPINES, MARIGOLD, MORNING GLORY, NASTURTIUM, PERIWINKLE, PETUNIA, PHLOX, PORTULACA, RUDBECKIA, SALVIA, SCABIOSA, SHASTA DAISY, SNAPDRAGON, STOCKS, SNOW ON THE MOUNTAIN, SUNFLOWER, SWEET ALYSSUM, SWEET PEA, SWEET WILLIAM, ZINNIA.

DO NOT USE ON GROUND COVER AS INJURY MAY RESULT.

**DIRECTIONS FOR MIXING INTO SOIL:** Apply Garden Weed Preventer to ornamentals established trees, flowers, and roses that are 2-3 inches or more in height. Apply uniformly at the recommended rate. The rate of application varies with the soil type. Use the table below to determine the proper rate for your type of soil. Mix into the soil surface with a hand rake, being care-

BEST DOCUMENT AVAILABLE

**DIRECTIONS FOR MIXING INTO SOIL:** Apply Garden Weed Preventer to ornamentals, established trees, flowers, and roses that are 2-3 inches or more in height. Apply uniformly at the recommended rate. The rate of application varies with the soil type. Use the table below to determine the proper rate for your type of soil. Mix into the soil surface with a hand rake, being careful not to disturb the roots of established plants. Shallow cultivation during the growing season will not reduce the herbicides effectiveness when it has been adequately mixed into the soil. Remove all existing grasses and weeds prior to application since weeds are controlled from seed, only.

**VEGETABLES:** Apply Garden Weed Preventer thoroughly and evenly at the recommended rate over the entire area. For large areas, use a seeder or broadcast applicator. Immediately after application mix into the soil to a depth of 1-3 inches using a rotary tiller or hand rake. Mixing should be done in two directions, at right angles, to insure thorough mixing.

**SPECIFIC VEGETABLE DIRECTIONS:**

From seed; CARROTS, OKRA, SOUTHERN PEAS, FIELD PEAS and BLACK-EYED PEAS: Apply anytime up to 10 weeks before planting from seed at the proper rate according to your soil type.

From seed; BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER, TOMATOES;

Apply anytime up to 10 weeks before planting, however use these rates:

✓ In light to medium soil, 1 lb. per 1280 sq. ft. and in heavy soil, 1 lb. per 960 sq. ft.

✓ Direct-seeded cole crops have exhibited marginal tolerance to recommended rates of Trifluralin. Stunting or reduced stands may occur.

Transplants; BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER, PEPPERS, TOMATOES: Apply and mix into the soil prior to transplanting. Do not apply after transplanting. See application rates below, for your type of soil.

IRISH POTATOES: Apply and mix into the soil after planting potatoes.

CUCUMBERS, CANTALOPES: Apply to dry foliage when plants have 3-4 true leaves. Mix thoroughly into the soil at the proper rate.

WEEDS CONTROLLED

**ANNUAL BROADLEAF WEEDS:** carelessweed, carpetweed, chickweed, florida, purslane, goosefoot, knotweed, kochia, Lambsquarters, pigweed, Russian thistle, stinging nettle

**ANNUAL GRASSES:** annual bluegrass, barnyardgrass, brachiaria, bromegrass, cheat, crabgrasses, foxtails, goosegrass, johnsongrass(from seed), junglerice, sandbur, stinkgrass, sprangletop, texas panicum, wildcane.

\*\*Gladiolus corms less than 1" in diameter may be injured.

NOTE: When applying Garden Weed Preventer to all established plants be sure that foliage is dry. Avoid overapplication since Garden Weed Preventer may cause injury to crops sensitive to Trifluralin planted the following season.

APPLICATION RATES WHEN MIXING GARDEN WEED PREVENTER INTO SOIL

SOIL TEXTURE	LIGHT	MEDIUM	HEAVY
	Sandy & Sandy Loam	Loam	Silt & Clay
APPLICATION RATES	1 lb.per 1280 sq.ft.	1 lb.per 960 sq.ft.	1 lb.per 640 sq.ft.

**Seacoast**  
LABORATORIES, INC.  
EAST BRUNSWICK, N.J. 08816

BEST DOCUMENT AVAILABLE