

GALLONS

NON-WARRANTY

This product is sold as is, without warranty of any kind, express or implied, including but not limited to the fitness for a particular purpose. The buyer and user assume all liability for any damage or injury resulting from the use of this product. The seller and manufacturer disclaim any liability for such damage or injury.

No agent or representative of the seller or manufacturer is authorized to make any representations or promises of warranty in connection with the sale of this product.



B-40

If the buyer and user do not accept this material upon these terms and conditions, it is to be returned, unopened, to the place of purchase within seven days. The use of this material shall be conclusive evidence that the buyer and user accept and agree to all the above-stated terms and conditions.

CAUTION! MAY CAUSE SKIN IRRITATION

Avoid Contact with Eyes, Skin, and Clothing.
Harmful if swallowed.

NOTE: When used as directed, this product is safe. It may be necessary to wear protective clothing and shoes when using this product.

Sold by
MIDLAND COOPERATIVES, INCORPORATED
MINNEAPOLIS, MINNESOTA 55413

2,4-D WEED KILLER

2,4-D EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENT
INERT INGREDIENTS

EPA Registration No. 1180-15

CAUTION: KEEP OUT OF REACH OF CHILDREN
See side panels for additional cautions

59

MIDLAND B-40 2/4-D
CONTROLS MANY BROAD-LEAVED WEEDS
AND CERTAIN WOODY PERENNIALS

- Canada thistle
- Cocklebur
- Dandelion
- Field bindweed
- Frenchweed
- Plantain

Hagweed 24
 Sumac 24
 Sow thistle 16
 Sunflower 24
 Wild morning 24
 Wild mustard 24

DIRECTIONS

The following recommendations for weed destruction in crops are intended to weed kill under good growing conditions in the early stage of weed growth

[illegible]

WHEAT AND BARLEY: * Use 1/2 to 1 pint per acre in sufficient water for uniform coverage. Spray in spring while weeds are young and grain is past jointing or heading.

Early, but stage. Do not spray grain in early seedling stage, nor during cotton from root
to root stage to milk stage. Fall spraying of winter wheat is not recommended. Use low
concentrations only in the weeds, such as Plantain, Common Yellow Mustard, and
other weeds. Conditions are favorable

OATS: 4 lb. or less when 2.4 ft. from other grains and should be sprayed in the "progs" at least 10-12 inches and 10-12 inches and before jointing time. put per acre. Do not apply if rain expected in 24 hr. or longer.

[illegible]

NON-SELECTIVE WEED CONTROL - *Lactuca* spp. appearing in a field.

$$\begin{aligned}
\mathcal{L}(\mathbf{y}|\mathbf{X}) &= \prod_{i=1}^n \frac{1}{\sigma_i} \exp\left\{-\frac{1}{2\sigma_i^2}(\mathbf{y}_i - \mathbf{X}_i^T \boldsymbol{\beta})^2\right\} \\
&= \frac{1}{\prod_{i=1}^n \sigma_i} \exp\left\{-\frac{1}{2} \sum_{i=1}^n \frac{(\mathbf{y}_i - \mathbf{X}_i^T \boldsymbol{\beta})^2}{\sigma_i^2}\right\} \\
&= \frac{1}{\prod_{i=1}^n \sigma_i} \exp\left\{-\frac{1}{2} \sum_{i=1}^n \frac{(\mathbf{y}_i - \mathbf{X}_i^T \boldsymbol{\beta})^2}{\sigma_i^2}\right\} \\
&= \frac{1}{\prod_{i=1}^n \sigma_i} \exp\left\{-\frac{1}{2} \sum_{i=1}^n \frac{(\mathbf{y}_i - \mathbf{X}_i^T \boldsymbol{\beta})^2}{\sigma_i^2}\right\}
\end{aligned}$$

MIDLAND COOPERATIVES, INCORPORATED
MINNEAPOLIS, MINNESOTA 55413