

Without any conditions beyond their control, neither Midland Cooperatives Inc. hereinafter called the Manufacturer, nor any intervening broker, distributor, dealer or agent, makes any representation or warranty, expressed or implied, with respect to results or safety from the use or handling of the above-mentioned material, and the buyer and user of this material assume all risk of loss, damage or injury which may result from the use or handling of this material, unless it is used in accordance with directions on this label.

No employee, salesman or other representative of the Manufacturer is authorized to alter or vary the terms or conditions herein stated, nor to make any representations or promises at variance with such terms and conditions.

If the buyer and user do not accept this material upon these terms and conditions, it is to be returned, unopened, to the place of purchase within seven days. The use of this material shall be conclusive evidence that the buyer and user accept and agree to all the above-stated terms and conditions.

**CAUTION: MAY CAUSE SKIN IRRITATION**  
**Avoid Contact with Eyes, Skin and Clothing**  
**HARMFUL IF SWALLOWED**

April 12,  
1973  
UNDER THE FEDERAL  
AGING SIDE AND ROD  
FOR ECONOMIC POIS  
UNDER NO. 1186 - 12  
13

Sold by  
**MIDLAND COOPERATIVES, INCORPORATED**  
MINNEAPOLIS, MINNESOTA 55421

# MCP WEED KILLER

**FOR WHEAT, BARLEY, OATS, RYE AND FLAX**

Contains 4 Pounds of OTC Acid Equivalent per Gallon at 4.0%		
<b>ACTIVE INGREDIENTS:</b>	Formic Acid (Active Ingredient) Methyl 4-Chlorophenoxyacrylate A-1	<b>BY WEIGHT</b>
<b>INERT INGREDIENTS</b>	Total	<b>100.00%</b>
Equivalent to Methyl 4-Chlorophenoxyacrylate A-1 4.0%		
<b>EPA Registration No. 1180-18</b>		

**CAUTION:** KEEP OUT OF REACH OF CHILDREN  
See side panels for additional cautions

General  
 Counsel  
 Office  
 1000  
 Pennsylvania  
 Avenue, N.W.  
 Washington, D.C. 20540  
 Tel: (202) 696-6000  
 Fax: (202) 696-6001  
 E-mail: [general\\_counsel@hhs.gov](mailto:general_counsel@hhs.gov)

**PREPARATION OF THE SPRAY** Fill the tank with hot water, required amount, then add the recommended amount of **MIDLAND MCP WEED KILLER** with agitation. Continue filling the spray tank with water enough water per acre to give uniform age. The amount of water required for volume applications may vary from 50 gallons per acre. For high volume applications 25 to 100 gallons, or more, of water needed for good coverage. In any case the same amount of **MIDLAND MCP KILLER** recommended per acre.

## CAUTION

Although MIDLAND MCP WEED KILLER volatile, care must be used to prevent

drift from coming into direct contact with tables, flowers, grapes, fruit trees, ornamental cotton, or other desirable plants sensitive to D. Coarse sprays are less likely to

DO NOT contaminate irrigation ditches or  
used for domestic purposes.

DO NOT use the same spray equipment for applying other materials to plants as may result.

DO NOT store near fertilizers, seed insecticides or fungicides.

DO NOT reuse container. Destroy when  
30 and 55 gallon drums may be returned  
reconditioning

CAUTION: MAY CAUSE SKIN IRRITATION  
Avoid Contact with Eyes, Skin and Clothing  
HARMFUL IF SWALLOWED

Sold by  
**MIDLAND COOPERATIVES, INCORPORATED**  
MINNEAPOLIS, MINNESOTA 55422

ONS

®

LLER

AND FLAX

162 F

BY WEIGHT

OF CHILDREN  
ditional cautions

## GENERAL INFORMATION

**PREPARATION OF THE SPRAY:** Fill the spray tank with half the required amount of water, then add the recommended amount of MIDLAND MCP WEED KILLER with agitation and continue filling the spray tank with water. Use enough water per acre to give uniform coverage. The amount of water required for low volume applications may vary from 5 to 25 gallons per acre. For high volume applications 25 to 100 gallons, or more, of water will be needed for good coverage. In any case, use the same amount of MIDLAND MCP WEED KILLER recommended per acre.

## CAUTION

Although MIDLAND MCP WEED KILLER is non-volatile, care must be used to prevent spray

drift from coming into direct contact with vegetables, flowers, grapes, fruit trees, ornamental cotton, or other desirable plants sensitive to 2, 4 D. Coarse sprays are less likely to drift.

DO NOT contaminate irrigation ditches or water used for domestic purposes.

DO NOT use the same spray equipment for applying other materials to plants, as injury may result.

DO NOT store near fertilizers, seed, insecticides, or fungicides.

DO NOT reuse container. Destroy when empty. 30 and 55 gallon drums may be returned for reconditioning.

**CAUTION: MAY CAUSE SKIN IRRITATION**  
Avoid Contact with Eyes, Skin and Clothing  
HARMFUL IF SWALLOWED

Sold by  
**MIDLAND COOPERATIVES, INCORPORATED**  
MINNEAPOLIS, MINNESOTA 55421

## MIDLAND MCP WEED KILLER CONTROLS THESE WEEDS

**SUSCEPTIBLE** (most areas): Lambsquarters, Mustard (annual), Ragweed, Stink weed, Wild Radish, Yellow Rocket or Wintercress, Cocklebur, Goatsbeard, Kochia, Shepherd's Purse and many others.

**LESS SUSCEPTIBLE:** Buttercup, Canada Thistle, Dandelion, Plantain, Hempnettle, Stinging Nettle, Sowthistle, Vetch, Sunflower, Pigweed and many others.

For the greatest effectiveness of MIDLAND MCP WEED KILLER apply when the weeds are young and vigorously growing.

## DIRECTIONS FOR USE

**WHEAT, BARLEY, OATS, RYE:** Apply as a water mix spray by ground sprayer or airplane. Use  $\frac{1}{2}$  to 1 pint per acre after crop has reached the 3 to 4 leaf stage up to boot stage. Do not spray from boot to dough stage.

**SMALL GRAINS UNDERSEEDED WITH ALFALFA, BIRDSFOOT TREFOIL, RED AND WHITE CLOVER:** For control of mustard, yellow rocket and other susceptible broadleaf weeds, apply  $\frac{1}{4}$  to  $\frac{1}{2}$  pint in not more than 5 to 10 gallons of water per acre. Make application after grain is well tillered (4 to 8 inches tall) and before reaching boot stage. The nurse crop and weeds should provide a protective canopy, which together with the use of low gallonage applied at low pressure will reduce the risk of damage to the legumes. Do not apply to small grains underseeded with vetch or sweet clover, which are very susceptible. There is also some risk to other legumes if only thinly protected by a canopy.

**FLAX:** Use  $\frac{1}{4}$  to  $\frac{1}{2}$  pint per acre. Apply only when weeds are up and when flax is 4 to 8 inches high and before it comes into bud. Treatment after early bud stage may result in severe damage. If Canada thistle is present it may be necessary to go as high as  $\frac{3}{4}$  pint per acre to prevent seed head production. Some injury to the flax may result.

**NOTE:** Regulations governing airplane applications of herbicides are in effect in many states. Consult local regulatory agencies concerning requirements.