



## RABON™ FLY-CON PREMIX

### RABON 7.76 ORAL LARVICIDE PREMIX

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	7.76%
INERT INGREDIENTS (consisting of Wheat Middlings)	92.24%
TOTAL	100.00%

### KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional precautionary statements.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS

### CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

When fed to cattle, RABON® Oral Larvicide passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch. By preventing larval development RABON® Oral Larvicide helps to control horn flies and face flies on pasture and house flies and stable flies in and around cattle lots.

MANUFACTURED BY  
MIDLAND COOPERATIVES, INCORPORATED  
Minneapolis, Minnesota


NET WEIGHT 50 LBS.



RABON® is a registered trademark of Shell Chemical Company  
U.S. Patent No. 3,192,842

EPA Reg. No. 1180-45 / Made in USA

PRODUCT CODE 50581-1-1/76



**RABON** Fly-Con Premix will mix uniformly in cattle feeds when standard mixing procedures are followed. Thus usual problems that are common to all feed preparation and which cause stratification, such as excessive free-fall or excessive handling, are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution of **RABON** Fly-Con Premix throughout the feed mix. The premix should be preblended with ground grain, protein supplements, mineral mixes, etc., before being added to roughages, such as chopped hay or silage. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if **RABON** Fly-Con Premix is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixers, horizontal blenders, mixer-feeder truck) may be used to prepare formulated feeds.

Feeds prepared with **RABON** Fly-Con Premix should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, **RABON** Fly-Con Premix should not be mixed in liquid feed supplements.

#### IN A CONCENTRATE FEED - ROUGHAGE FED SEPARATELY

Midland **RABON** Fly-Con Premix can be used to prepare concentrate feeds that will provide 792 mg. of **RABON** per animal per day.

To prepare a larvicidal concentrate feed, mix Midland **RABON** Fly-Con Premix according to the amount of concentrate to be fed per animal per day. Use the following table as a guide for determining the proper mixing rate.

#### MIXING DIRECTIONS

Lbs. of Concentrate Consumed per Animal Per Day	<b>RABON</b> in the Concentrate mg./lb.	Lbs. of <b>RABON</b> Fly-Con Premix per Ton of Concentrate
.5	1584	0.35
1.0	792	0.18
1.5	528	0.12
2.0	396	0.087
5.0	159	0.035
10.0	79	0.018
15.0	53	0.012
20.0	39	0.0087
25.0	32	0.0069
30.0	26	0.0059

Feed the appropriate larvicidal concentrate to cattle weighing between 400 and 1,200 lbs. For larger cattle weighing between 1,200-1,700 lbs., increase the amount of concentrate fed to 1 1/2 times that recommended.

Start feeding **RABON** larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

Rations containing **RABON** Fly-Con Premix may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.

**RABON** Fly-Con Premix prevents the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle but is not effective against existing adult flies. In some cases, supplemental fly control measures may be needed in and around cattle lots and barns to control adult house flies and stable flies which can breed not only in cattle manure but other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, **RABON** Fly-Con Premix should be used in conjunction with other good management and sanitation practices.

#### IN A COMPLETE RATION - NO OTHER ROUGHAGE FED

**RABON** Fly-Con Premix can be used to prepare rations that contain 26.4 mg. of **RABON** per lb. of complete ration.

To prepare a larvicidal ration, mix 1.5 lbs. of **RABON** Fly-Con Premix per ton of complete mixed ration containing both grain and roughage.

Full-feed this larvicidal complete ration to feeder cattle weighing from 400 to 1,400 lbs. or to dairy cattle at a rate to sustain milk production, but not less than 2.6 lbs. of the ration per 100 lbs. of body weight daily.

#### STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or by other approved methods for pesticide containers, or bury in a sanitary place.

NOTED