

TREATMENT: Administer the agent to the patient. Antiprimitive is the drug against parasympathetic nervous system. If there are signs of bradycardia, atropine sulfate should be injected at 1 to 2 mg intervals. A dose of 1 to 2 mg. grams of the complete atropinization has been reported.

- Morphine is contraindicated • Monitor for urinary retention • Observe patient continuously for 4-6 hours • Repeat drug administration as necessary • Patients may without warning cause a hypotensive episode by taking the drug with any other decrease inhibitor • Allow no further drug administration until the drug is completely metabolized • Has been allowed as determined by the state.

DIRECTIONS FOR APPLYING **LETHALAIRE** A-30

IMPORTANT — First read the precautions printed on Right Panel of Label
ENTOMOLOGICAL RECOMMENDATIONS —
 Lasioderma serricorne
GENERAL INFORMATION LETHALAIRE A 30

PREPARATION OF TOBACCO FACTORY AND WAREHOUSE AREAS

PREPARATION OF NON FOOD FACTORY OR SEED WAREHOUSE AREAS

LETHALAIRE A 30

FLYING INSECTS

LETHALAIRE A 30

STORED PRODUCT AND CRAWLING INSECTS

LETHALAIRE A 30

NON AIR CONDITIONED FACTORY AREAS

LETHALAIRE A 30

AIR CONDITIONED FACTORY AREAS

LETHALAIRE A 30

LETHALAIRE A 30

LOT
NO.

UN 1956

DOT SHIPPING NAME
COMPRESSED GAS, n o s

LETHALAIRE

AN AEROSOL INSECTICIDE CONTAINING

20% DDVP

FOR THE CONTROL OF TOBACCO INSECTS.

ACTIVE INGREDIENTS

INERT INGREDIENTS



**KEEP OUT OF REACH OF CHILDREN
 VAPOR EXTREMELY HAZARDOUS
 DANGER! POISON**

SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

ALWAYS USE AN APPROVED GAS MASK WHEN APPLYING

PHYSICAL OR CHEMICAL HAZARDS

THIS CYLINDER IS REFILLABLE BY MANUFACTURER
 Return to Manufacturer for Deposit or Exchange

EPA EST 1187-VA 1

EPA Reg No

VIRGINIA CHEMICALS INC. PORTSMOUTH, VA

ETHALAIRE

IL INSECTICIDE CONTAINING

0% DDVP

CONTROL OF TOBACCO INSECTS.

AND CRAWLING INSECTS.
WHERE FERTILIZERS ARE USED
IN THE FIELD, KEEP AWAY FROM

A-30

NET WT.

15 LBS.

IS

**KEEP OUT OF REACH OF CHILDREN
VAPOR EXTREMELY HAZARDOUS
DANGER! POISON**

RIGHT PANEL FOR ADDITIONAL PRECAUTIONS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Do not breathe vapor. If inhaled, move to fresh air. If swallowed, do not induce vomiting. If on skin, wash with soap and water. If in eyes, flush with water. Keep away from food and feed. Do not eat or drink from treated areas.

IMPROVED GAS MASK WHEN APPLYING - When applying, wear a gas mask with a canister. Do not eat or drink from treated areas. Do not breathe vapor. If inhaled, move to fresh air. If swallowed, do not induce vomiting. If on skin, wash with soap and water. If in eyes, flush with water. Keep away from food and feed. Do not eat or drink from treated areas.

PHYSICAL OR CHEMICAL HAZARDS

Do not breathe vapor. If inhaled, move to fresh air. If swallowed, do not induce vomiting. If on skin, wash with soap and water. If in eyes, flush with water. Keep away from food and feed. Do not eat or drink from treated areas.

THIS CYLINDER IS REFILLABLE BY MANUFACTURER
Turn to Manufacturer for Deposit or Exchange when Empty

EPA EST 1187-VA-1

EPA Reg No 1187-67

ETHALAIRE CHEMICALS INC. PORTSMOUTH, VIRGINIA

STORAGE AND DISPOSAL

Do not store in areas where food or feed is stored. Do not store in areas where children or domestic animals are present. Do not store in areas where food or feed is stored. Do not store in areas where children or domestic animals are present.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Do not breathe vapor. If inhaled, move to fresh air. If swallowed, do not induce vomiting. If on skin, wash with soap and water. If in eyes, flush with water. Keep away from food and feed. Do not eat or drink from treated areas.

FIRST AID TREATMENT

Do not breathe vapor. If inhaled, move to fresh air. If swallowed, do not induce vomiting. If on skin, wash with soap and water. If in eyes, flush with water. Keep away from food and feed. Do not eat or drink from treated areas.

NOTE TO PHYSICIAN

Do not breathe vapor. If inhaled, move to fresh air. If swallowed, do not induce vomiting. If on skin, wash with soap and water. If in eyes, flush with water. Keep away from food and feed. Do not eat or drink from treated areas.

