

Sold to Professional Pest Control Operators Only

DIRECTIONS: To be used by licensed pest control operators only. Use one 12 oz. can for each unobstructed closed area of 10,000 cubic feet but not less than 7,000 cubic feet or one 6 oz. can for each 5,000 cubic feet. During fogging and to release contents, use a mask or respirator of a type approved by the U.S. Bureau of Mines for DDVP insecticide protection. Open cabinets and doors to areas to be treated. Shut off air conditioners and fans before using.

All occupants must be removed from house or building before treatment is started and should not reoccupy building for at least 2 hours. Open doors and windows and allow to be ventilated before occupying. Unwrapped or unpeached foods should be removed or placed in refrigerator during fogging. Food handling equipment, dishes, should be washed after application. Do not contaminate water, food or foodstuffs. Remove all pets. Cover fish tanks, or remove. Cover house plants within 8 feet of point of release with newspaper to prevent possible burning of tender plants.

Place canister on table in center of area to be treated, or near point of greatest infestation. Place newspapers under canister. Use one or more canisters in combinations of 6 and 12 ounce sizes to give the best distribution of vapors where free flow of mist is not assured.

Snap lock-open nozzle over tip of container. Leave building at once. The entire contents will release automatically.

WARNINGS: Clean Out is poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or eyes. Do not breathe fumes. Wear natural rubber gloves, protective clothing and goggles when handling. Wear a mask or respirator of a type approved by the U.S. Bureau of Mines for DDVP VAPOR insecticide protection. Keep all unprotected persons out of operational area. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before smoking or eating. Wash all contaminated clothing with soap and hot water before reuse. Contents are under pressure. High temperature may burst container. Do not expose to direct sunlight, radiators, stove, hot water or other sources of heat.



27 228 973

NET WT. 6 OUNCES
(170 GRAMS)



Clean-Out

AEROSOL
INSECTICIDE

with 7% VAPONA* Insecticide and Ronnel

**KILLS EXPOSED STAGES OF ROACHES,
TICKS, FLEAS, FLIES AND SILVERFISH
IN HOMES AND BUILDINGS**

ACTIVE INGREDIENTS: 2,2-dichlorovinyl dimethyl phosphate (DDVP) 6.5%, related compounds 0.5%, Ronnel (0,0-dimethyl 0-(2, 4, 5-trichlorophenyl) phosphorothioate 2.00%, Petroleum Distillates 4.00%, Xylol 2.00%.

INERT INGREDIENTS: 85.00%
Equivalent to 7% VAPONA Insecticide

DANGER:

KEEP OUT OF REACH OF CHILDREN



POISON



See antidote statement
and other cautions on side panel.

E.P.A. Reg. No. 1187-80

VIRGINIA CHEMICALS INC. *Shell
PORTSMOUTH, VA. 23703 Reg. T.M.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE is the emergency antidote for VAPONA insecticide poisoning. Consult your physician about an adequate supply of 1.00 grain atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoonful of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1.00 grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See warning symptoms below.) KEEP patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN:

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minute intervals in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may without warning cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.


PRECAUTIONS IN USING: Clean Out is poisonous, therefore precautions must be observed when handling. (See antidote and warnings.) Store in a safe place. Children or animals must not be able to reach or handle. Care should be used in handling empty containers. Do not contaminate food, water or foodstuffs.

DISPOSAL OF CONTAINERS: Completely empty container by removing container to an open air area away from pets and animals. Press release button until no more vapor escapes. Puncture can to prevent reuse. Bury can 18 inches deep in an isolated location away from water supplies.

ACCEPTED
JAN 7 1974
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO.

1187-80

Comments



Cameraman's Note
Poor Copy

ACCEPTED WITH COMMENTS