

DIRECTIONS FOR APPLICATION

LETHALAIRE A-35

LETHALA

RECOMMENDATIONS: LETHALAIRE A-35 is a contact insecticide. It is effective against a wide range of tobacco insects, including the tobacco budworm, tobacco rootworm, and tobacco beetle. It is also effective against the tobacco moth and the tobacco fly. LETHALAIRE A-35 is a contact insecticide. It is effective against a wide range of tobacco insects, including the tobacco budworm, tobacco rootworm, and tobacco beetle. It is also effective against the tobacco moth and the tobacco fly.

GENERAL INFORMATION: LETHALAIRE A-35 is a contact insecticide. It is effective against a wide range of tobacco insects, including the tobacco budworm, tobacco rootworm, and tobacco beetle. It is also effective against the tobacco moth and the tobacco fly. LETHALAIRE A-35 is a contact insecticide. It is effective against a wide range of tobacco insects, including the tobacco budworm, tobacco rootworm, and tobacco beetle. It is also effective against the tobacco moth and the tobacco fly.

CAUTION: LETHALAIRE A-35 is a contact insecticide. It is effective against a wide range of tobacco insects, including the tobacco budworm, tobacco rootworm, and tobacco beetle. It is also effective against the tobacco moth and the tobacco fly. LETHALAIRE A-35 is a contact insecticide. It is effective against a wide range of tobacco insects, including the tobacco budworm, tobacco rootworm, and tobacco beetle. It is also effective against the tobacco moth and the tobacco fly.

AN AEROSOL INSECTICIDE CONTAINING
40% VAPONA
FOR THE CONTROL OF TOBACCO INSECTS

Virginia Chemicals

BEFORE USING THIS PRODUCT READ THESE PRECAUTIONS

POISON



CAUTION



POISON

- 1. DO NOT CONTACT THE SKIN.** Gloves impervious to DDVP should be worn. All skin surfaces should be completely covered with protective clothing during application.
- 2. DO NOT INHALE DDVP MIST.** Wear a gas mask of a type approved by the U. S. Bureau of Mines for DDVP protection.
- 3. ADEQUATE PERSONAL HYGIENE AND CLEANLINESS** are necessary. After applying, remove clothing and bathe thoroughly with warm water and soap. Wash clothing before re-use.
- 4. DO NOT CONTAMINATE FOOD,** smoke, eat or chew tobacco in the operating area.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE is the emergency antidote for DDVP insecticide poisoning. 2-PAM is also antidotal and may be administered in conjunction with atropine. Consult your physician about obtaining an adequate supply of 1/100-grain atropine tablets for emergency use. **CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100-grain atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** (See WARNING SYMPTOMS below).

- Keep patient prone and quiet.
- Start artificial respiration immediately if patient is not breathing.
- Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred.

- Morphine is contraindicated.
- Clear chest by postural drainage.
- Observe patient continuously for 48 hours.
- Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor.
- Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

	By Weight
.....	37.2%
.....	2.8%
.....	60.0%
.....	100.0%

ISON

CHILDREN

ARDOUS



0. Poisonous if swallowed, label carefully. Use only as directed. Do not breathe vapor. Do not get solid or liquid in eyes. Keep out of reach of children. Wash off clothing.

IN APPLYING and have an
Bureau of Mines for DDVP
ergency.

No. 1187-108

als Inc.

'03